

THE EISENHOWER LIBRARY



3 1151 02690 5921

1880

Library



Johns Hopkins of the University



THE EFFECT OF ONE SALT
UPON
THE HYDRATING POWER OF ANOTHER SALT
PRESENT IN THE SAME SOLUTION.

DISSERTATION

Submitted to the Board of University Studies of the
Johns Hopkins University in Conformity with
the Requirements for the Degree of
Doctor of Philosophy

By

Charles M. A. Stine.

1907.

THE NEW YORK PUBLIC LIBRARY

ASTOR LENOX TILDEN FOUNDATION

1215 Broadway, New York, N. Y.

1850

1215 Broadway, New York, N. Y.

THE NEW YORK PUBLIC LIBRARY

ASTOR LENOX TILDEN FOUNDATION

1215 Broadway, New York, N. Y.

1215 Broadway, New York, N. Y.

1215

1215 Broadway, New York, N. Y.

1215

CONTENTS.

Acknowledgment - - - - -	3
Introduction - - - - -	4
Conductivity Calculations - - - - -	8
Experimental:	
Freezing-Point - - - - -	-14
Conductivity - - - - -	16
Measurements:	
Calcium Chloride - - - - -	19, 44
Potassium Chloride - - - - -	25, 141
Mixture Of Calcium Chloride And Potassium Chloride -	27
Magnesium Chloride - - - - -	46
Mixture Of Calcium Chloride And Magnesium Chloride -	48
Strontium Chloride - - - - -	62
Mixture Of Calcium Chloride And Strontium Chloride -	64
Magnesium Nitrate - - - - -	-79, 107
Strontium Nitrate - - - - -	-81
Mixture Of Magnesium Nitrate And Strontium Nitrate -	83
Sodium Bromide - - - - -	91
Lithium Bromide - - - - -	-93
Mixture Of Lithium Bromide And Sodium Bromide - - -	-95
Calcium Nitrate - - - - -	105

1. The first step in the process of identifying a problem is to define the problem. This involves identifying the symptoms of the problem and determining the scope of the problem. Once the problem has been defined, the next step is to identify the causes of the problem. This involves identifying the factors that are contributing to the problem and determining the underlying causes of the problem. Once the causes of the problem have been identified, the next step is to develop a plan of action to address the problem. This involves identifying the steps that need to be taken to solve the problem and determining the resources that will be needed to implement the plan.

2. The second step in the process of identifying a problem is to identify the causes of the problem. This involves identifying the factors that are contributing to the problem and determining the underlying causes of the problem. Once the causes of the problem have been identified, the next step is to develop a plan of action to address the problem. This involves identifying the steps that need to be taken to solve the problem and determining the resources that will be needed to implement the plan.

3. The third step in the process of identifying a problem is to develop a plan of action to address the problem. This involves identifying the steps that need to be taken to solve the problem and determining the resources that will be needed to implement the plan. Once the plan of action has been developed, the next step is to implement the plan. This involves carrying out the steps that have been identified in the plan of action and monitoring the progress of the implementation.

4. The fourth step in the process of identifying a problem is to implement the plan. This involves carrying out the steps that have been identified in the plan of action and monitoring the progress of the implementation. Once the plan has been implemented, the next step is to evaluate the results of the implementation. This involves assessing the effectiveness of the plan and determining whether the problem has been solved.

5. The fifth step in the process of identifying a problem is to evaluate the results of the implementation. This involves assessing the effectiveness of the plan and determining whether the problem has been solved. Once the results of the implementation have been evaluated, the next step is to identify the lessons learned from the process. This involves identifying the factors that contributed to the success or failure of the implementation and determining the steps that need to be taken to improve the process.

6. The sixth step in the process of identifying a problem is to identify the lessons learned from the process. This involves identifying the factors that contributed to the success or failure of the implementation and determining the steps that need to be taken to improve the process. Once the lessons learned have been identified, the next step is to develop a plan of action to address the lessons learned. This involves identifying the steps that need to be taken to improve the process and determining the resources that will be needed to implement the plan.

7. The seventh step in the process of identifying a problem is to develop a plan of action to address the lessons learned. This involves identifying the steps that need to be taken to improve the process and determining the resources that will be needed to implement the plan. Once the plan of action has been developed, the next step is to implement the plan. This involves carrying out the steps that have been identified in the plan of action and monitoring the progress of the implementation.

Mixture Of Calcium Nitrate And Magnesium Nitrate - -	109
Aluminium Chloride - - - - -	118
Ferric Chloride - - - - -	120
Mixture Of Ferric Chloride And Aluminium Chloride -	122
Mixture Of Calcium Nitrate And Calcium Chloride - -	132
The Effect Of Change In Temperature Upon The Conductivities Of Separate Solutions Of Electrolytes And Upon Mixtures Of These Electrolytes - - - - -	140
Discussion Of Results - - - - -	147
Summary - - - - -	149
Biography - - - - -	151

ACKNOWLEDGMENT.

To President Remsen, Professor Morse and Professor Jones, the author takes pleasure in expressing his gratitude for instruction both in the laboratory and lecture room.

He desires also to thank Dr. Tingle, Dr. Swartz and Dr. Acree for instruction and advice received from them.

This investigation was undertaken at the suggestion of Professor Jones, and was carried out under his supervision.

1
THE EFFECT OF ONE SALT
UPON
THE HYDRATING POWER OF ANOTHER SALT
PRESENT IN THE SAME SOLUTION.

Eight years ago Jones and Ota¹ and Jones and Knight² observed that certain salts in concentrated solution, give abnormally great lowering of the freezing-point of water. The problem was taken up experimentally by Jones and Chambers³ and studied for a number of chlorides and bromides. As a result of this work Jones⁴ concluded that "In concentrated solutions these chlorides and bromides must take up part of the water, forming complex compounds with it, and thus removing it from the field of action as far as freezing-point lowering is concerned. The compound, which is probably very unstable, formed by the union of a molecule of the chloride or bromide with a large number of molecules of water, acts as a unit or as one molecule in lowering the freezing-point of the remaining water. The total amount of water present, which is now acting as solvent, is diminished by the amount taken up by the chloride or bromide molecules. The lowering of the freezing-point is thus abnormally great, because a part of the water is no longer present as solvent, but is

1. Amer. Chem. Journ., 22, 5 (1899).

2. Ibid., 22, 110 (1899).

3. Ibid., 23, 89 (1900).

4. Ibid., 23, 103 (1900).

In combination with the chloride or bromide molecules."

A large amount of work on this problem has since been carried on in this laboratory and nineteen papers¹ have already been published by Jones and his coworkers in this laboratory on the combination of solvent and dissolved substance. We shall not attempt to review the work already done, but shall merely state briefly the lines of evidence developed for the hydrate conception, and then proceed to the consideration of our particular phase of the problem.

Five distinct lines of evidence bearing upon the hydrate theory have been pointed out: First, the minima in the boiling-point curves manifest themselves at greater concentrations than in the freezing-point curves. "This is just

-
1. Jones and Chambers: Amer. Chem. Journ., 23, 169 (1900).
 - Chambers and Frazer: Ibid., 23, 512 (1900).
 - Jones and Getman: Ibid., 27, 433 (1902).
 - " " " : Ztschr. phys. chem., 46, 244 (1903); Phys. Rev., 18, 146 (1904).
 - Jones and Getman: Amer. Chem. Journ., 31, 303 (1904).
 - " " " : Ztschr. phys. chem., 49, 365 (1904).
 - " " " : Ber. d. chem. Ges., 37, 1511 (1904).
 - " " " : Amer. Chem. Journ., 32, 308 (1904).
 - " " " : Ibid., 32, 338 (1904).
 - Jones and Bassett: Amer. Chem. Journ., 33, 534 (1905).
 - " " " : " " " 34, 390 (1905); J. chim. Phys., 6, 455 (1905).
 - Jones and Bassett: Z. phys. chem., 52, 231 (1905).
 - " " " " 55, 365 (1906).
 - Jones and McMaster: Amer. Chem. Journ., 35, 186 (1906).
 - " " " " 35, 445 (1906).
 - " " " : Monograph #60, Carnegie Inst.

what would be expected in terms of our hydrate theory, since, at the higher temperature, the hydrates would be less stable and a greater concentration would have to be reached before there would be sufficient hydration to change the direction of the curve and produce a minimum."

Second; the relation between water of crystallization and lowering of freezing-point.¹ Hydrating power is a function of water of crystallization—the larger the number of molecules of water of crystallization the greater, in general, is the hydrating power of the salt. The water of crystallization may be taken as roughly proportional to the hydrating power of the substance.

Third; the complexity of the hydrates, as well as the number of molecules of water of crystallization varies with the temperature?

Fourth; certain color changes² which take place in solution, e.g., the change in color of a solution of cobalt chloride from a typical purplish red to a beautiful blue when treated with a small amount of some other salt with large hydrating power, such as calcium chloride or aluminum chloride. This effect of water of hydration upon the

1. Amer. Chem. Journ., 31, 344 (1904); Ztschr. phys. chem., 49, 385 (1904); Amer. Chem. Journ., 32, 308 (1904); 33, 534 (1905); J. chim. Phys., 3, 455 (1905).

2. Amer. Chem. Journ., 34, 293 (1905).

3. " " " " , 34, 290 (1905).

vibration of the dissolved particles is shown by the spectrograms of various binary mixtures of salts.¹

Fifth; the temperature coefficients of conductivity² of aqueous solutions of electrolytes are greater, the greater the hydrating power of the electrolyte. The temperature coefficients of conductivity of aqueous solutions of electrolytes are of the same order of magnitude for those solutions having, approximately, the same hydrating power. The temperature coefficients of conductivity for any given substance increase with the dilution of the solution, and this increase is greatest for those substances with large hydrating power, just as would be expected from the hydrate theory.

Assuming, then, the existence of hydrates to be sufficiently well established by the work referred to above, it seemed desirable to attempt to determine, if possible, the effect of one salt with hydrating power upon the hydrate formed by a second salt in the same solution. Also the effect upon the hydrates formed by a salt, when there is present in the same solution a second salt with very small hydrating power. When both salts possess the power of forming

1. Amer. Chem. Journ., 29 (1907); Monograph #60, Carnegie Inst.

2. Amer. Chem. Journ., 28, 445 (1906).

hydrates in single solution, the amounts of water taken up by the salts at the same concentrations may be the same, or vary within wide limits. We proposed to study binary mixtures in which both conditions are illustrated.

Since we desired to calculate the approximate composition of the hydrates formed in the mixtures, as nearly as possible, it was necessary to obtain data for the freezing-points, conductivities, and number of grams of solvent in a litre of solution of the various salts employed. In order that we might have the data for the exact normalities of the solutions which we mixed, it seemed advisable to make the necessary measurements in the single solutions, rather than to attempt to interpolate from data previously obtained.

CONDUCTIVITY CALCULATIONS.

It is well known that when two or more electrolytes are present in the same solution, the conductivity of this solution is not equal to the sum of the conductivities which these electrolytes have in separate solutions of the same normality as that which these salts may be assumed to have in the mixture; but that the conductivity of the mixture is generally less than this sum. If the solutions are isohydric¹, that is, the component solutions possess a common

1. Arrhenius: Wied. Ann., 30, 51 (1857); Ztschr. phys. Chem., 2, 281 (1886).

ion in equal concentration, then the simplest conditions prevail between the conductivities of the single solution and that of the mixture. This latter condition exists when

$$\frac{\alpha_1}{V_1} = \frac{\alpha_2}{V_2}$$

in which α_1 is the dissociation of one of the electrolytes at the dilution V_1 , and α_2 is the dissociation of the other electrolyte at the dilution V_2 . Such solutions can be mixed in any proportion without altering their dissociation, and hence the conductivity of the mixture is the mean of the conductivities of the component solutions, and can, therefore, be calculated according to the following formula:

$$k = \frac{k_1 P + k_2 P'}{P + P'}$$

in which k_1 and k_2 are the conductivities, and P and P' the proportions of the component solutions.

Unfortunately, we have to deal with no such simple relations as this in the mixtures with which we worked. If both electrolytes have a common ion and each is a weak electrolyte, then they would obey the law of mass action. Wakeran¹ and Larmwater² have approached this case from the standpoint of isohydricism.

Wolf³ has investigated the conductivity of mixtures

1. Ztschr. phys. chem., 15, 159 (1894).

2. " " " , 45, 557 (1903).

the first of these is the fact that the
 second of these is the fact that the
 third of these is the fact that the

$$\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$$

the first of these is the fact that the
 second of these is the fact that the
 third of these is the fact that the
 fourth of these is the fact that the
 fifth of these is the fact that the
 sixth of these is the fact that the
 seventh of these is the fact that the
 eighth of these is the fact that the

$$\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$$

the first of these is the fact that the
 second of these is the fact that the
 third of these is the fact that the
 fourth of these is the fact that the
 fifth of these is the fact that the
 sixth of these is the fact that the
 seventh of these is the fact that the
 eighth of these is the fact that the

the first of these is the fact that the
 second of these is the fact that the
 third of these is the fact that the
 fourth of these is the fact that the
 fifth of these is the fact that the
 sixth of these is the fact that the
 seventh of these is the fact that the
 eighth of these is the fact that the

of dilute solutions of strong electrolytes, with solutions of weak electrolytes. Selrader⁴ has calculated the degree of dissociation of each electrolyte, from considerations of the transport numbers of the ions, the quantities of the anions and the cations migrating to their respective poles, the quantities of each separated at the electrodes, etc. Hoffmeister⁵ and Hopfgartner⁶ have dealt with the concentrations of the cations in a mixture of two electrolytes. Novas and Koro⁷ indicate a method of ascertaining the dissociation of the salts in a mixed solution, which they apply to the solutions with which they had to deal. Barnwater's⁸ equations take into consideration the viscosity of the solution, the friction between the ions and the solvent, and between the ions themselves, the normality, dissociation, current employed, etc., and give a satisfactory agreement between calculated and experimental results. MacGreor⁹ and his students have also carried out considerable

1. Ztschr. Electrochem. 3, 498 (1897).
5. " phys. chem., 27, 345 (1898).
6. " " 25, 115 (1898).
7. Journ. Chem. Soc., 24, 1148 (1902).
8. Ztschr. phys. chem., 23, 426 (1899).
9. Trans. N. S. Inst. of Sci., 10, (1898 - 99); 9, 101 (1895 - 96); Phil. Mag., 41, 276 (1896); Trans. Roy. Soc. Can., 2, sect. 3, 15 (1899 - 97).

work upon the conductivities of mixed electrolytes, solving the equations deduced by a graphical method.

In our work, however, the viscosity of the mixtures differs considerably, in many cases, from that of the single solutions assumed to have the same normality. This would involve changes in friction in the movement of the ions. Also, the amount of water combined as hydrate, i. e., the dimensions of the atmosphere about the ion, continually changed from one normality to another. This involved a varying degree of freedom of movement of the ions. Again, the variation in the number of particles present, ions and molecules, must involve continual variation in the amount of friction between the ions. As a comparatively short, and at the same time sufficiently accurate method, we have adopted the following in apportioning the diminution in conductivity to the constituent salts:— Since we have dealt with mixtures in which the two electrolytes have a common ion in every case, we have based the method on the number of this common ion yielded by the constituent salts in separate solutions at the same normality. By the same normality we mean that if equal volumes of two solutions are mixed, and any diminution in volume which may take place

upon mixing, is compensated for by the addition of pure solvent, then the normality of each solution in the mixture may be assumed to be exactly half of that of the constituent solutions. If suppression of the ionization is dependent upon the common ion, then it will be inversely as the number of this common ion furnished by each of the salts in a mixture. This number was obtained thus:— The figure representing the normality of the individual solutions, multiplied by the value of α for the solution at the normality in question, would give the gramme atomic weight of the common ion. Suppose the case to be one where a solution of potassium chloride has been combined with an equal volume of a solution of calcium chloride. It being assumed that calcium chloride dissociates into



this figure representing the gramme atomic weight of the common ion would have to be doubled for a ternary electrolyte, such as calcium chloride. Since two chlorine ions are required to re-form a molecule of calcium chloride, the figure expressing the number of chlorine ions driven back in the case of calcium chloride must be divided by two, in order to get the relative number of molecules of calcium chloride re-formed.

Let us take the values of Λ_{∞} as expressing the molecular conductivity of the completely dissociated molecules of the various salts. If the figures for the relative numbers of molecules of the two salts re-formed, be multiplied by the proper value of Λ_{∞} , we shall then have the relative losses in conductivity. From this the absolute losses are readily obtained. An example may suffice to make this clear:- Take the case of the mixture of equal volumes of a solution of 1.8 N. calcium chloride and of 1.7 N. potassium chloride. The resulting solution, any diminution of volume having been compensated for, will consist of 0.9 N. calcium chloride and 0.85 N. potassium chloride. The dissociation of a 0.9 N. solution of the specimen of calcium chloride employed was found by conductivity measurements to be 53.2%; of 0.85 N. potassium chloride, 83.2%. Doubling the value of α in the case of calcium chloride and multiplying by the normality, we get 0.9576. Similarly, for potassium chloride, 0.7068. These represent the relative numbers of chlorine ions which would be furnished by each salt in the mixture, if no suppression of the ionization took place upon mixing. That is, if driving back occurs, according to the law of mass action, 707 chlorine ions

should be driven back to re-form calcium chloride; to 15% for potassium chloride. But since it requires two ions of chlorine to re-form each molecule of calcium chloride, only 45% molecules of calcium chloride will be re-formed to every 95% molecules of potassium chloride. Multiplying these values by Λ_{∞} for calcium chloride and potassium chloride respectively, we get 5323, approximately, for calcium chloride; and 7695 for potassium chloride. These values represent the relative losses in conductivity upon mixing. From this the decrease in conductivity can be apportioned between the two salts.

All conductivity measurements are given as specific conductivities in the mixtures. The units employed are reciprocal centimetre ohms. For purposes of comparison with the previous work upon the "Hydrate theory" our units may be transformed by dividing Λ_v by 1.0×9.1 . This would give μ_v for the same liquid.

The compositions of the hydrates in the single solutions was calculated by the method previously employed by Jones² and his coworkers.

EXPERIMENTAL WORK.

FREEZING POINT. The more concentrated solutions were frozen by means of a mixture of solid carbon dioxide and alcohol, the freezing temperature being determined by means of alcohol thermometers. These temperatures could not be measured more closely than 0.5. For the more dilute solu-

ions freezing mixtures of ice and sodium chloride, or ice and crystallized calcium chloride were employed. The temperatures were read by means of Beckmann thermometers, covering a range of 6° , 12° and 25° respectively. When the more concentrated solutions were frozen they were removed from the bath of solid carbon dioxide and alcohol as soon as they began to separate, in order that they might not be surrounded by the very cold freezing mixture while temperature equilibrium was being established. The necessary correction for the separation of ice and consequent increase in concentration was introduced in every instance. The separation of some of the dissolved substance was readily ascertained by the grit or sand like character of the solid that separated, which was easily detected. Since it was found that in the volume of liquid employed in the freezing-point tube, about twenty-five cubic centimeters, a freezing-point determination made with a large undercooling, and consequent copious separation of ice, gave somewhat different results from one where the undercooling was less, care was taken

-
1. "The Conductivity of Liquids," O. F. Tower, pages 2 et seq.
 2. Amer. Chem. Journ., 34, 292 (1905); Amer. Chem. Journ., 35, 545 (1905).

both in the freezing-point determinations of the pure solvent and of the solutions, to keep the undercooling nearly the same:—about a degree, throughout. This necessitated inoculating the solution with an exceedingly minute particle of the solid phase of the solvent, in nearly every case.

CONDUCTIVITY.

Conductivities were measured by the method of Kohlrausch, employing a Wheatstone bridge, telephone receiver and induction coil. The bridge wire was of "Manganin" and was calibrated before beginning the work, by the method of Strouhal and Marcus.¹

The resistance coils, manufactured by Leeds and Co., were tested and found in no case to involve a greater error than 0.04 per cent.

The dilute solutions were measured in conductivity cells of the form employed by Jones and McJaster;² the tubes carrying the platinum electrodes being sealed into both the upper and lower walls of the stopper. For the more concentrated solutions cells of the type used by Jones and Getman³

1. *Wied. Ann.*, 10, 326 (1880).

2. *Amer. Chem. Journ.*

3. *Itzchr. phys. Chem.*, 46, 254 (1903).

were employed. The electrodes were treated in the usual manner (coated with platinum black, etc.), in order to obtain a sharper tone minimum or point of silence in the readings.

The cells for the more dilute solutions were standardized by determining the cell constant by means of $N/50$ potassium chloride. The value $k = .0015183$ was taken as the conductivity of $N/50$ potassium chloride. For the more concentrated solutions the cell constants were determined with a $N/2$ solution of potassium chloride, for which $k = .08365$ at 0° .⁶ The potassium chloride employed in determining cell constants was recrystallized five times, the last two times from conductivity water, and was free from all impurities. The cell constants were frequently redetermined, but very little change in them was noted from time to time. The cells were filled with distilled water when not in use, and were always carefully cleansed, dried, and rinsed with the new solution when a change in solutions was made. All conductivities were measured at 0° . The temperature of the zero bath was always within a tenth of a degree of 0° as determined by an accurately standardized thermom-

⁶These values are calculated from the results of Jones and West. See Amer. Chem. Journ., 54, 381 (1905).

eter. All conductivity measurements are the average of four measurements with differing resistances in circuit.

The water used in this work was purified thus: Ordinary distilled water was redistilled from potassium dichromate and sulphuric acid. This was then again distilled from a second flask containing acidified potassium dichromate, and the steam led through a boiling solution of barium hydroxide and condensed in a block tin condenser. The water purified as above had a conductivity of 1.2 to 2.5 $\times 10^{-6}$ at 0°.

It is well known that the conductivity method is not an accurate measure of dissociation in concentrated solutions. Especially is this the case at the concentrations with which we dealt, since in the mixtures we have employed, the values of Λ_{∞} were determined for each salt by diluting the single solution to a maximum value of Λ_{ν} , as the value which Λ_{∞} would exhibit in the mixture. This is, obviously, a source of additional error, for the conductivity of the ions resulting from a completely dissociated molecule in a very dilute solution must be different from the conductivity of these ions in a more concentrated solution. In the more dilute solution a larger atmosphere of the solvent is attached to the ions, and this must be dragged along by

them. There is a much smaller number of particles of salt (ions or molecules), present in the dilute solution than in the more concentrated. Consequently the friction between the ions in their movements must be different. The viscosity of the solutions in which Λ_{∞} can be determined is very much less than the viscosity of the solutions at the concentrations employed. These are all important factors which affect the conductivity of any dissolved substance. The method is, however, an approximate one for measuring such dissociations and is the best we can do at present. All that can be said is, that it probably gives values of the right order of magnitude.

CALCIUM CHLORIDE.

A strong mother solution of Calcium chloride was made up and standardized by precipitating the calcium in a known volume, and from this calculating the amount of calcium chloride present. All solutions were made up by diluting the calculated amount of this mother solution to the required volume.

Table I. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ/m .
0.3	1.517	5.056
0.5	2.673	5.344
0.6	3.270	5.450
0.7	4.065	5.807
0.9	5.422	6.024
1.0	6.410	6.410
1.1	7.080	6.450
1.35	9.362	6.934
1.4	10.05	7.18
1.55	11.58	7.47
1.755	14.33	8.16
1.88	14.69	8.16
2.3	21.07	9.58
2.7	30.25	11.20
3.1	39.5	12.74
3.5106	49.5 ?	14.13 ?

In this table, as elsewhere where they occur, m is the concentration in terms of gram molecules per liter; Δ the observed freezing-point lowering, corrected for the separation of ice, and Δ/m the molecular lowering of the freezing-point.

Table II. CONDUCTIVITY MEASUREMENTS.

$$\Lambda_{\infty}, 0^{\circ} = 147.52.1$$

m.	v.	k.	Λ_m .	α .
0.2	3.533	.028044	98.5	.674
0.5	2.000	.045701	67.0	.594
0.6	1.6666	.050266	63.8	.568
0.7	1.4285	.057555	62.2	.557
0.9	1.1111	.070601	78.4	.532
1.0	1.000	.07554	75.5	.512
1.1	0.909	.081573	74.1	.502
1.35	0.7407	.092860	68.6	.466
1.4	0.7143	.09396	67.1	.455
1.55	0.6453	.099184	64.0	.434
1.755	0.5698	.10733	61.1	.414
1.8	0.5555	.107367	59.8	.405
2.2	0.4545	.115676	53.7	.357
2.7	0.3704	.127757	45.6	.300
3.1	0.3226	.11410	36.8	.250
3.5106	0.2848	.10710	30.5	.207

In this table, as elsewhere throughout this paper, m is the concentration, in terms of gram-molecules per liter; v is the number of liters that contain a gram-molecule of the dissolved substance; k is the specific conductivity,

1. Calculated from the value given by Jones and Dassel; Amer. Chem. Journ., 23, 507 (1905).

and is equal to

$$\frac{\mu_{\alpha} \times .001009}{v} ;$$

Λ_v is the molecular conductivity at 0° , and is equal to $1000 \times v \times k$; α is the per cent of dissociation, and is equal to $\Lambda_v / \Lambda_{\infty}$.

Table III. WEIGHT-NORMAL CORRECTIONS, or Number of Grams of Solvent in a 1000 cubic centimeters of Solution.

N.	W Sol.	W Salt	W H ₂ O	Correction Per Cent.
0.3	51.270	1.685	49.045	0.8
0.5	52.2608	2.775	49.2988	1.0
0.6	52.7437	3.33	49.4137	1.2
0.7	53.1162	3.885	49.3512	1.5
0.9	53.9816	4.995	48.9866	2.0
1.0	54.346	5.55	48.776	2.4
1.1	54.772	6.105	48.667	2.7
1.25	55.9435	7.4925	48.451	3.1
1.4	56.115	7.770	48.345	3.3
1.55	56.6562	8.6025	48.0537	3.9
1.755	57.7040	9.7402	47.9638	4.1
1.8	28.900	4.995	23.905	4.1
2.2	29.650	6.105	23.545	5.8
2.7	30.882	7.492	23.390	7.3
3.1	31.461	8.602	22.859	8.6
3.5106	32.187	9.720	22.467	10.1

In these tables of weight-normal corrections, m has the usual significance; $W_{\text{Sol.}}$ is the weight of twenty-five or of fifty cubic centimeters of the solution; W_{Salt} is the weight of salt contained in this volume of the solution; $W_{\text{H}_2\text{O}}$ is the weight of water in this volume of solution, equal to $W_{\text{Sol.}} - W_{\text{Salt}}$; Correction Per Cent is the percentage correction to be applied in order to refer measurements to a thousand grams of solvent. These are not specific gravity measurements made in pycnometers, and we therefore give the correction per cent only to the first decimal.

In the following table, as in all other tables of the hydrates formed in the single solutions, m has the usual significance; α is the per cent of dissociation; L is the theoretical molecular lowering of the freezing-point referred to a thousand grams of the solvent; Δ/m the molecular lowering found experimentally; L' the molecular lowering, n the number of gram-molecules of water in combination, both being referred to a thousand grams of solvent;

$$M = \frac{1000 - \frac{L}{L'} \times 1000}{18} ;$$

M is the number of molecules of water in combination with one molecule of the salt, at the concentration in question, if a liter of the solution at that concentration, contained a thousand grams of water: it equals $\frac{M}{m}$.

Table IV.

HYDRATES.

m.	α .	L.	Δ/m .	L ¹ .	M.	H.
0.3	.674	4.22	5.06	5.02	8.6	29.4
0.5	.594	4.00	5.34	5.29	12.6	25.2
0.6	.568	3.97	5.45	5.29	14.6	24.3
0.7	.557	3.93	5.81	5.72	17.4	24.8
0.9	.532	3.84	6.02	5.90	19.4	21.6
1.0	.512	3.76	6.31	6.16	21.6	"
1.1	.502	3.73	6.44	6.26	22.4	20.4
1.35	.466	3.59	6.93	6.72	25.9	19.2
1.4	.455	3.55	7.18	6.94	27.2	19.4
1.55	.434	3.47	7.47	7.18	28.7	16.5
1.755	.414	3.40	8.16	7.83	31.4	17.9
1.8	.405	3.37	"	7.80	31.6	17.5
2.2	.357	3.19	9.58	9.03	35.9	16.3
2.7	.296	2.96	11.20	10.39	39.7	14.7
3.1	.259	2.79	12.74	11.65	43.6	14.1
3.5106	.207	2.63	14.10	12.68	46.5	13.2

Since we wished to investigate the effect of the presence of a salt with little hydrating power upon a salt with considerable hydrating power, potassium chloride was added to calcium chloride. The data for the various solutions of potassium chloride alone are therefore given next.

No.	Date	Part	Description		Total	
			Debit	Credit	Debit	Credit
100	10/1	100.00	100.00		100.00	
101	10/2	100.00	100.00		100.00	
102	10/3	100.00	100.00		100.00	
103	10/4	100.00	100.00		100.00	
104	10/5	100.00	100.00		100.00	
105	10/6	100.00	100.00		100.00	
106	10/7	100.00	100.00		100.00	
107	10/8	100.00	100.00		100.00	
108	10/9	100.00	100.00		100.00	
109	10/10	100.00	100.00		100.00	
110	10/11	100.00	100.00		100.00	
111	10/12	100.00	100.00		100.00	
112	10/13	100.00	100.00		100.00	
113	10/14	100.00	100.00		100.00	
114	10/15	100.00	100.00		100.00	
115	10/16	100.00	100.00		100.00	
116	10/17	100.00	100.00		100.00	
117	10/18	100.00	100.00		100.00	
118	10/19	100.00	100.00		100.00	
119	10/20	100.00	100.00		100.00	
120	10/21	100.00	100.00		100.00	
121	10/22	100.00	100.00		100.00	
122	10/23	100.00	100.00		100.00	
123	10/24	100.00	100.00		100.00	
124	10/25	100.00	100.00		100.00	
125	10/26	100.00	100.00		100.00	
126	10/27	100.00	100.00		100.00	
127	10/28	100.00	100.00		100.00	
128	10/29	100.00	100.00		100.00	
129	10/30	100.00	100.00		100.00	
130	10/31	100.00	100.00		100.00	

These are the only entries in the ledger for the month of October.

The total of the debit column is \$10,000.00 and the total of the credit column is \$10,000.00.

The difference between the debit and credit columns is \$0.00.

The balance of the account is \$10,000.00.

The total of the debit column is \$10,000.00 and the total of the credit column is \$10,000.00.

POTASSIUM CHLORIDE.

Table V. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ/m .
0.45	1.952	3.955
0.65	2.24	3.45
0.85	2.91	3.43
.90	3.08	3.42
1.05	3.58	3.41
1.2972	4.46	3.45
1.3	4.49	3.46
1.7	5.93	3.49
2.1	7.43	3.54
2.5945	9.29	3.58

Table VI. CONDUCTIVITY MEASUREMENTS.₁
 $\Lambda_{\infty}, 0^{\circ} = 80.33.$

m.	v.	k.	Λ	α .
0.45	2.223	.030382	67.4	.839
0.65	1.538	.043608	67.1	.835
0.85	1.1765	.050804	66.6	.832
0.90	1.1111	.053717	65.2	.819
1.05	0.9523	.061969	65.7	.816
1.2972	0.7709	.087948	64.7	.805
1.3	0.7392	.084118	"	"
1.7	0.5882	.109202	64.2	.800
2.1	0.4762	.133785	63.7	.793
2.5945	0.3851	.162207	62.8	.782

Table VII. WEIGHT-NORMAL CORRECTIONS.

m.	"Sol.	Wsalt.	W _{H₂O} .	Correction PER CENT.
0.45	6.0042	0.20353	7.74007	1.4
0.65	6.0735	0.38066	7.69284	2.3
0.85	6.1447	0.49654	7.64766	2.6
0.90	25.9986	1.6785	24.3151	2.7
1.05	6.2169	0.61492	7.60198	2.2
1.3	26.5127	2.4245	24.0882	3.7
1.7	26.928	3.1705	23.7575	5.0
2.1	27.4457	3.9165	23.5292	5.9
2.5945	28.0254	4.8385	23.1869	7.3

Table VIII. HYDRATES.

m.	α .	L.	Δ/m .	L ¹ .	M.	H.
0.45	.829	3.42	3.975	3.80		
0.65	.835	3.41	3.45	3.78		
0.85	.832	"	3.43	3.74		
0.90	.812	3.37	3.42	3.32		
1.05	.818	3.38	3.41	3.40		
1.2972.605		3.56	3.45	3.53		
1.3	"	"	3.46	"		
1.7	.800	3.35	3.49	3.31		
2.1	.793	3.34	3.54	3.33		
2.5945.762		3.32	3.58	3.32		

Evidently no hydrating power.

1. Calculated from the value given by Jones and West; Amer. Chem. Journ., 37, 381 (1905).

Table 1: Summary of the data sets

Dataset	N	N_{train}	N_{test}	N_{val}
1	10000	8000	1000	1000
2	10000	8000	1000	1000
3	10000	8000	1000	1000
4	10000	8000	1000	1000
5	10000	8000	1000	1000
6	10000	8000	1000	1000
7	10000	8000	1000	1000
8	10000	8000	1000	1000
9	10000	8000	1000	1000
10	10000	8000	1000	1000

Table 2: Summary of the data sets

Dataset	N	N_{train}	N_{test}	N_{val}
1	10000	8000	1000	1000
2	10000	8000	1000	1000
3	10000	8000	1000	1000
4	10000	8000	1000	1000
5	10000	8000	1000	1000
6	10000	8000	1000	1000
7	10000	8000	1000	1000
8	10000	8000	1000	1000
9	10000	8000	1000	1000
10	10000	8000	1000	1000

Table 1 and Table 2 show the summary of the data sets.

MIXTURE OF CALCIUM CHLORIDE AND POTASSIUM CHLORIDE.

Table IX. FREEZING-POINT MEASUREMENTS.

m.	P.	Δ .
1.0 <chem>CaCl2</chem>	1	
0.9 K Cl	"	4.413
1.4 <chem>CaCl2</chem>	1	
1.3 K Cl	"	6.625
1.8 <chem>CaCl2</chem>	1	
1.7 K Cl	"	9.17
2.2 <chem>CaCl2</chem>	1	
2.1 K Cl	"	12.01
2.7 <chem>CaCl2</chem>	1	
2.5945 K Cl	"	15.00
5.1 <chem>CaCl2</chem>	1	
2.5945 K Cl	"	16.53
3.5196 <chem>CaCl2</chem>	1	
2.5945 K Cl	"	18.0

Column headed P indicates parts by volume of each solution which were mixed; m is the normality of the solutions before mixing; In order that each solution should have exactly one-half the normality after mixing which it had before, the volume of the mixture was brought to exactly twice the volume of either constituent solution.

Table 1. Summary of the results of the analysis of variance for the different parameters of the model.

Parameter	Source of variation	Sum of squares
Mean	1	1.0000
Standard deviation	1	1.0000
Skewness	1	1.0000
Kurtosis	1	1.0000
Mean	1	1.0000
Standard deviation	1	1.0000
Skewness	1	1.0000
Kurtosis	1	1.0000
Mean	1	1.0000
Standard deviation	1	1.0000
Skewness	1	1.0000
Kurtosis	1	1.0000

The results of the analysis of variance for the different parameters of the model are presented in Table 1. The results show that the mean, standard deviation, skewness and kurtosis of the data are all significantly different from zero. This indicates that the data are not normally distributed. The results also show that the mean, standard deviation, skewness and kurtosis of the data are all significantly different from each other. This indicates that the data are not normally distributed.

Table 1. Summary of the results of the analysis of variance for the different parameters of the model.

TABLE X. CONDUCTIVITY MEASUREMENTS.

ρ .	ρ .	π_{eq} .	v_{eq} .	k_{eq} .	k .	D .	κ_{eq} .	Δ_{eq} .	α .
1.0 CaCl_2	1	0.5	2.000	.089077	.048801	.004156	.04880	94.6	.572
0.9 CaCl_2	"	0.45	2.222		.030352		.027607	61.5	.736
1.4 CaCl_2	1	0.7	1.438		.057555		.054352	72.1	.593
1.3 CaCl_2	"	0.65	1.538	.038713	.048908	.00745	.039087	80.0	.742
1.3 CaCl_2	1	0.9	1.111		.070601		.065221	75.5	.401
1.7 CaCl_2	"	0.95	1.173	.114096	.056804	.013309	.048875	52.5	.716
2.0 CaCl_2	1	1.1	0.909		.031573		.072052	62.7	.452
2.1 CaCl_2	"	1.05	0.952	.13081	.063063	.01373	.057453	54.7	.621
2.7 CaCl_2	1	1.55	0.7407		.032360		.033341	53.6	.401
2.5345 CaCl_2	"	1.2372	0.7703	.14305	.033343	.02273	.06765	52.0	.642
2.1 CaCl_2	1	1.55	0.8451		.030194	.03252	.087734	55.4	.375
2.5345 CaCl_2	"	1.2372	0.7703	.15031	.033040		.065012	50.1	.654
2.5103 CaCl_2	1	1.755	0.5203	.152337	.07028		.02193	52.3	.355
2.5345 CaCl_2	"	1.2372	0.7703		.033040	.03279	.06053	46.7	.531

N is the normality of the solutions mixed. P is the parts by volume of each solution which were mixed. When equal volumes are mixed the resulting normality of each solution is exactly one-half of that of the solutions before mixing. This is given under n_0 . V_0 is number of liters of solution which will contain a gram molecular weight of the salt, based on n_0 . K_u is the specific conductivity of the mixture. This would be the sum of the conductivities of the component solutions if no suppression of the ionization occurred. K gives the conductivities of the single solutions of each salt at the normality acquired upon mixing. In each case the sum of these two, minus the conductivity of the mixture, k_u , gives the decrease in conductivity due to the driving back of the dissociation of the salts in question. This is given under D . This loss in conductivity is then apportioned to each electrolyte by the method already described, and the resulting conductivities are given under K_0 . From this, Λ_v and α are obtained.

Table XI. WEIGHT PERCENT COMPOSITION.

M.	P.	WSol.	WSalt.	WH ₂ O.	Correction Per Cent
1.0 CaCl	1	26.7148	2.22667	24.51843	1.9
0.9 KCl	"				
1.4 CaCl	1	27.5662	3.1542	24.3120	5.2
1.3 KCl	"				
1.8 CaCl	1	28.0354	4.0827	23.9527	4.2
1.7 KCl	"				
2.2 CaCl	1	28.6177	5.0102	23.6075	5.6
2.1 KCl	"				
2.7 CaCl	1	29.422	6.1655	23.2565	6.9
2.5945 KCl	"				
3.1 CaCl	1	29.8247	6.7205	23.1042	7.6
2.5945 KCl	"				
3.5106 CaCl	1	30.2118	7.2753	22.9235	8.3
2.5945 KCl	"				

Proceeding upon the tentative hypothesis that the calcium chloride forms the same hydrates in the presence of a salt with no hydrating power, that it forms in separate solution, Tables XII and XIII are prepared.

No. of molecules	Molecular Weight			No. of atoms
	$\text{C}_{10}\text{H}_{16}\text{O}_4$	$\text{C}_{10}\text{H}_{16}\text{O}_3$	$\text{C}_{10}\text{H}_{16}\text{O}_2$	
1.0	100.00	100.00	100.00	1.0000
1.5	150.00	150.00	150.00	1.5000
2.0	200.00	200.00	200.00	2.0000
2.5	250.00	250.00	250.00	2.5000
3.0	300.00	300.00	300.00	3.0000
3.5	350.00	350.00	350.00	3.5000
4.0	400.00	400.00	400.00	4.0000
4.5	450.00	450.00	450.00	4.5000
5.0	500.00	500.00	500.00	5.0000

For each molecule, the molecular weight is calculated by summing the atomic weights of all the atoms in the molecule. The molecular weight is then divided by the number of atoms in the molecule to give the average molecular weight. This is the value used in the calculations.

Table XII. CALCIUM CHLORIDE IN THE MIXTURE.

m.	P.	m _c .	α .	L.	I.
1.0 CaCl ₂ 0.9 KCl	1 "	0.5	.573	3.992	2.031
1.4 CaCl ₂ 1.3 KCl	1 "	0.7	.599	3.829	2.769
1.8 CaCl ₂ 1.7 KCl	1 "	0.9	.691	3.687	3.161
3.2 CaCl ₂ 3.1 KCl	1 "	1.1	.452	3.541	4.126
2.7 CaCl ₂ 2.5945 KCl	1 "	1.35	.404	3.362	4.88
3.1 CaCl ₂ 2.5945 KCl	1 "	1.55	.375	3.256	5.162
3.5106 CaCl ₂ 2.5945 KCl	1 "	1.7553	.355	3.179	6.084

α is the dissociation of the calcium chloride in the mixture. L is the freezing-point lowering calculated from the formula $(1.86 \times 2\alpha) + 1.86$. I is the freezing-point lowering, corrected to one thousand grams of solvent, which the salt would give if there were no hydration.

m.	P.	m_c .	α .	Δ .	W_{H_2O}	Correction Per Cent	I.
1.0 CaCl 0.9 KCl	$\frac{1}{2}$ "	0.45	.766	1.478	753.7	24.6	1.961
1.4 CaCl 1.3 KCl	$\frac{1}{2}$ "	0.65	.748	2.113	655.9	34.4	3.222
1.8 CaCl 1.7 KCl	$\frac{1}{2}$ "	0.85	.716	2.712	608.2	39.2	4.454
2.2 CaCl 2.1 KCl	$\frac{1}{2}$ "	1.05	.681	3.283	538.5	46.2	6.097
2.7 CaCl 2.5945 KCl	$\frac{1}{2}$ "	1.2972	.649	3.068	464.7	53.5	8.564
3.1 CaCl 2.5945 KCl	$\frac{1}{2}$ "	1.2972	.624	3.021	408.0	59.2	9.605
3.5106 CaCl 2.5945 KCl	$\frac{1}{2}$ "	1.2972	.581	3.941	340.5	66.0	11.20

α is the dissociation of the potassium chloride in the mixture. Δ is the freezing-point lowering deduced from $[(1.66 \times \alpha) + 1.86] m_c$. W_{H_2O} is the weight of water acting as solvent toward the potassium chloride, deduced from:—
 $L : L' :: x : 1000$, which gives the amount of water taken up by the calcium chloride in the formation of hydrates when it alone is present in solution, and the weight-normal correction for the mixture. I is the ideal lowering the potassium chloride would give if the calcium chloride formed the same hydrates in the mixture that it forms when alone in solution.

Table XIV. Comparison of lowerings due to hydration in separate solution of CALCIUM CHLORIDE and in the mixture.

m.	P.	m_c .	Δ_c .	I_c	Diff.	Δ .	I	Diff.
1.0 CaCl 0.9 KCl	$\frac{1}{11}$	0.5	4.413	3.995	0.418	3.673	3.066	0.606
1.1 CaCl 1.8 KCl	$\frac{1}{11}$	0.7	6.625	5.991	0.634	4.004	2.75	1.25
1.8 CaCl 1.7 KCl	$\frac{1}{11}$	0.9	9.170	7.918	1.252	5.314	3.455	1.859
2.3 CaCl 3.1 KCl	$\frac{1}{11}$	1.1	12.01	10.223	1.782	6.896	4.100	2.796
2.7 CaCl 2.5945 KCl	$\frac{1}{11}$	1.35	15.0	13.442	1.558	9.072	4.851	4.221
3.1 CaCl 2.5945 KCl	$\frac{1}{11}$	1.55	16.5	15.067	1.433	11.13	5.385	5.745
3.5106 CaCl 2.5945 KCl	$\frac{1}{11}$	1.7553	18.0	17.288	0.712	14.10	5.97	8.13

Δ_c is the lowering found experimentally for the mixture. I_c is the sum of I in Tables VII and XIII. $\text{Diff}_c = \Delta_c - I_c =$ lowering due to hydration. Δ is lowering found experimentally for calcium chloride, corrected to one thousand grams of solvent. I is the lowering the calcium chloride should have given in 1000 grams of solvent, if there were no hydration. Evidently Diff_c and Diff should be identical if there were no change in hydration. Such is not the case. Consequently, the basis assumed for the calculation of I for Potassium chloride in the mixture is incorrect, since it

depend upon a knowledge of the amount of water eliminated from the sphere of action as solvent, by the hydration of the calcium chloride. It, therefore, becomes necessary to derive a formula for the calculation of the hydration of the calcium chloride, in the mixture.

Formula for Calculating Lowering due to Hydration, in a Binary Mixture of Electrolytes, where One Salt has no Hydrating Power.

Let x° equal lowering due to hydration of the salt with hydrating power, in the mixture.

L = ideal lowering which the salt with hydrating power should give in the mixture in question, at the corrected dissociation, if there were no hydration.

v = number of litres which contain a gram molecule of the salt with hydrating power, at the normality it has in the mixture.

$$\text{Then } (x + 1)v = \Delta/m.$$

S = Correction to 1000 grams of solvent, necessary in the mixture.

$$(x + 1)v - (x + 1)vS = L'.$$

x' = amount of water acting as solvent toward the salt with no hydrating power.

$$L : L' :: x' : 1000.$$

$$L = (1.86 \times 2\alpha) + 1.86, \text{ in a ternary electrolyte,}$$

where α represents the corrected dissociation of the salt with hydrating power, in the mixture.

$$\frac{(1.86 \times 2\alpha) + 1.86}{(x + 1)v - (x + 1)v_3} \cdot 1000 = x'.$$

But taking into account the weight-normal correction also, in the mixture in question, the above becomes,

$$\left[\frac{(1.86 \times 2\alpha) + 1.86}{(x + 1)v - (x + 1)v_3} \cdot 1000 \right] - 108 = \text{amount of water act-}$$

ually acting as solvent toward salt with no hydrating power.

Putting this into the form of a percentage correction,

$$\frac{(1.86 \times 2\alpha) + 1.86}{(x + 1)v - (x + 1)v_3} - \frac{S}{100} = \text{the per cent by which}$$

I for the salt with little, or no hydrating power must be divided.

Let α' = the corrected dissociation of the salt with no hydrating power, in the mixture.

$[1.86 \times \alpha] + 1.86] m$ = lowering it would give in 1000 grams of solvent, if it is a binary electrolyte. m is the normality of the salt with no hydrating power.

$$\text{then, } \frac{1.86m(\alpha' + 1)}{(x + 1)(v - v_3)} - \frac{S}{100} + x + 1 = \text{lowering found}$$

experimentally for the mixture. From this expression the value of x may be derived.

By means of this formula the value of x has been cal-

The first two terms of the series are $\frac{1}{2}$ and $\frac{1}{4}$.

Since $\frac{1}{2} + \frac{1}{4} = \frac{3}{4}$, the sum of the first two terms is $\frac{3}{4}$.

$$\frac{1}{2} + \frac{1}{4} = \frac{2}{4} + \frac{1}{4} = \frac{3}{4}$$

The next two terms are $\frac{1}{8}$ and $\frac{1}{16}$. Their sum is $\frac{3}{16}$.

Adding this to the previous sum gives $\frac{3}{4} + \frac{3}{16} = \frac{12}{16} + \frac{3}{16} = \frac{15}{16}$.

$$\frac{3}{4} + \frac{3}{16} = \frac{12}{16} + \frac{3}{16} = \frac{15}{16}$$

The next two terms are $\frac{1}{32}$ and $\frac{1}{64}$. Their sum is $\frac{3}{64}$.

Adding this to the previous sum gives $\frac{15}{16} + \frac{3}{64} = \frac{150}{64} + \frac{3}{64} = \frac{153}{64}$.

$$\frac{15}{16} + \frac{3}{64} = \frac{150}{64} + \frac{3}{64} = \frac{153}{64}$$

The next two terms are $\frac{1}{128}$ and $\frac{1}{256}$. Their sum is $\frac{3}{256}$.

Adding this to the previous sum gives $\frac{153}{64} + \frac{3}{256} = \frac{1530}{256} + \frac{3}{256} = \frac{1533}{256}$.

The next two terms are $\frac{1}{512}$ and $\frac{1}{1024}$. Their sum is $\frac{3}{1024}$.

Adding this to the previous sum gives $\frac{1533}{256} + \frac{3}{1024} = \frac{15330}{1024} + \frac{3}{1024} = \frac{15333}{1024}$.

The next two terms are $\frac{1}{2048}$ and $\frac{1}{4096}$. Their sum is $\frac{3}{4096}$.

Adding this to the previous sum gives $\frac{15333}{1024} + \frac{3}{4096} = \frac{153330}{4096} + \frac{3}{4096} = \frac{153333}{4096}$.

The next two terms are $\frac{1}{16384}$ and $\frac{1}{32768}$. Their sum is $\frac{3}{32768}$.

$$\frac{153333}{4096} + \frac{3}{32768} = \frac{1533330}{32768} + \frac{3}{32768} = \frac{1533333}{32768}$$

The next two terms are $\frac{1}{131072}$ and $\frac{1}{262144}$. Their sum is $\frac{3}{262144}$.

Adding this to the previous sum gives $\frac{1533333}{32768} + \frac{3}{262144} = \frac{15333330}{262144} + \frac{3}{262144} = \frac{15333333}{262144}$.

The next two terms are $\frac{1}{1048576}$ and $\frac{1}{2097152}$. Their sum is $\frac{3}{2097152}$.

culated for the various concentrations employed in the mixture of calcium chloride and potassium chloride, and the results are given in Table IV. The significance of the abbreviations has already been given in connection with the preceding tables.

TABLE IV. HYDRATION OF CaCl_2 IN ALXPHT.

η .	P.	λ_c .	α .	L.	L.	I.	x.	Δ/π .	L!	L.	H.
1.0	CaCl_2	1	.572	3.92	2.924		0.453	4.391	4.396	10.2	15.3
0.2	KCl	"									
1.4	CaCl_2	1	.523	3.823	2.769		0.209	5.254	5.258	12.7	12.7
1.2	KCl	"									
1.2	CaCl_2	1	.431	3.637	3.464		1.527	5.558	5.558	17.1	10.0
1.7	KCl	"									
2.2	CaCl_2	1	.452	3.541	4.126		2.221	5.224	5.507	13.3	13.3
2.1	KCl	"									
2.7	CaCl_2	1	.404	3.332	4.33		2.34	5.722	5.337	20.2	15.5
2.5245	KCl	"									
2.1	CaCl_2	1	.375	3.256	5.432		3.601	5.247	5.478	22.1	14.2
2.5245	KCl	"									
2.5108	CaCl_2										
2.5245	KCl	"	.355	3.173	6.034		4.32	5.322	5.472	22.1	15.1


In order to compare the hydrates formed by calcium chloride in separate solution with those which it forms in the mixture with potassium chloride the values of N are plotted against the concentrations as abscissae, in Figure I.

It is seen that while the value of N increases with increase in concentration, in general, in the mixture as in separate solution, N is less in the mixture than in separate solution of calcium chloride. This is just what we should expect, since there is less water present as solvent in the mixture of calcium chloride and potassium chloride at any given concentration of the calcium chloride, than there is at the same concentration of the single solution of calcium chloride. Now it is evident from previous work that while the total number of gram-molecules of water combined increases in general, with increase in concentration, the molecular hydration decreases with increase in concentration. That is, the amount of water with which a molecule of the calcium chloride or the resulting ions combine, depends upon the number of molecules of water present to one molecule of the calcium chloride. If now, instead of increasing the number of grams of salt with hydrating power, present in a given solution, thus diminish-

I. CALCIUM CHLORIDE ALONE.

II. CALCIUM CHLORIDE MIXED WITH POTASSIUM CHLORIDE.

DEPT. EXPERIMENTAL ENGINEERING, SIKLEY COLLEGE, CORNELL UNIVERSITY.



GRAM-MOLECULES OF SALT PER LITER OF SOLUTION.

FIGURE I.

If the amount of water, we add to the calcium chloride a substance like potassium chloride and then make up to the former volume, it is evident that, there being less water present, the amount of water with which each molecule or ion of calcium chloride will combine will be diminished; that is, the value of M will be diminished.

In order to compare the difference between the amount of water present acting as solvent toward the calcium chloride, in the single solution and in the mixture, with the difference in the values of M in the two cases, we give in the following table, under D_{H_2O} the difference between the number of grams of water present as solvent in the separate solution of calcium chloride and in the mixture at the respective concentrations; under D_M the difference in the values of M for the separate solution of the calcium chloride and in the mixture at the same concentrations.

The first of these is the fact that the
 the second is the fact that the
 the third is the fact that the
 the fourth is the fact that the
 the fifth is the fact that the
 the sixth is the fact that the
 the seventh is the fact that the
 the eighth is the fact that the
 the ninth is the fact that the
 the tenth is the fact that the

The first of these is the fact that the
 the second is the fact that the
 the third is the fact that the
 the fourth is the fact that the
 the fifth is the fact that the
 the sixth is the fact that the
 the seventh is the fact that the
 the eighth is the fact that the
 the ninth is the fact that the
 the tenth is the fact that the
 the eleventh is the fact that the
 the twelfth is the fact that the
 the thirteenth is the fact that the
 the fourteenth is the fact that the
 the fifteenth is the fact that the
 the sixteenth is the fact that the
 the seventeenth is the fact that the
 the eighteenth is the fact that the
 the nineteenth is the fact that the
 the twentieth is the fact that the
 the twenty-first is the fact that the
 the twenty-second is the fact that the
 the twenty-third is the fact that the
 the twenty-fourth is the fact that the
 the twenty-fifth is the fact that the
 the twenty-sixth is the fact that the
 the twenty-seventh is the fact that the
 the twenty-eighth is the fact that the
 the twenty-ninth is the fact that the
 the thirtieth is the fact that the
 the thirty-first is the fact that the
 the thirty-second is the fact that the
 the thirty-third is the fact that the
 the thirty-fourth is the fact that the
 the thirty-fifth is the fact that the
 the thirty-sixth is the fact that the
 the thirty-seventh is the fact that the
 the thirty-eighth is the fact that the
 the thirty-ninth is the fact that the
 the fortieth is the fact that the
 the forty-first is the fact that the
 the forty-second is the fact that the
 the forty-third is the fact that the
 the forty-fourth is the fact that the
 the forty-fifth is the fact that the
 the forty-sixth is the fact that the
 the forty-seventh is the fact that the
 the forty-eighth is the fact that the
 the forty-ninth is the fact that the
 the fiftieth is the fact that the
 the fifty-first is the fact that the
 the fifty-second is the fact that the
 the fifty-third is the fact that the
 the fifty-fourth is the fact that the
 the fifty-fifth is the fact that the
 the fifty-sixth is the fact that the
 the fifty-seventh is the fact that the
 the fifty-eighth is the fact that the
 the fifty-ninth is the fact that the
 the sixtieth is the fact that the
 the sixty-first is the fact that the
 the sixty-second is the fact that the
 the sixty-third is the fact that the
 the sixty-fourth is the fact that the
 the sixty-fifth is the fact that the
 the sixty-sixth is the fact that the
 the sixty-seventh is the fact that the
 the sixty-eighth is the fact that the
 the sixty-ninth is the fact that the
 the seventieth is the fact that the
 the seventy-first is the fact that the
 the seventy-second is the fact that the
 the seventy-third is the fact that the
 the seventy-fourth is the fact that the
 the seventy-fifth is the fact that the
 the seventy-sixth is the fact that the
 the seventy-seventh is the fact that the
 the seventy-eighth is the fact that the
 the seventy-ninth is the fact that the
 the eightieth is the fact that the
 the eighty-first is the fact that the
 the eighty-second is the fact that the
 the eighty-third is the fact that the
 the eighty-fourth is the fact that the
 the eighty-fifth is the fact that the
 the eighty-sixth is the fact that the
 the eighty-seventh is the fact that the
 the eighty-eighth is the fact that the
 the eighty-ninth is the fact that the
 the ninetieth is the fact that the
 the ninety-first is the fact that the
 the ninety-second is the fact that the
 the ninety-third is the fact that the
 the ninety-fourth is the fact that the
 the ninety-fifth is the fact that the
 the ninety-sixth is the fact that the
 the ninety-seventh is the fact that the
 the ninety-eighth is the fact that the
 the ninety-ninth is the fact that the
 the hundredth is the fact that the

Table XVI.

m.	D_{H_2O} .	D_{KCl} .
0.5	9	2.4
0.7	17	4.8
0.9	22	2.34
1.1	29	2.61
1.25	38	4.99
1.55	" "	6.62
1.7558	42	8.36

These results are plotted in curves in Figure II.

For the first four concentrations the values of M decrease by the same amount, since the amount of potassium chloride added is increased as the calcium chloride is increased. Consequently, the curve for the difference in the values of M is approximately parallel to the axis of abscissas (concentration). Referring to Table XV it will be seen that the difference between successive values of M diminishes with increase in concentration, until finally in the mixture containing 2.7% calcium chloride and 2.5945 M potassium chloride, M has the value 20.9. The effect of the potassium chloride on the amount of water with which the calcium chloride can combine, is such that, although the

DIFFERENCE BETWEEN AMOUNT OF WATER PRESENT IN THE SINGLE SOLUTION OF CALCIUM CHLORIDE AND IN THE MIXTURE OF CALCIUM CHLORIDE AND POTASSIUM CHLORIDE
 . DIFFERENCE BETWEEN VALUES OF λ FOR CALCIUM CHLORIDE IN SINGLE SOLUTION OF CALCIUM CHLORIDE AND IN MIXTURE OF CALCIUM CHLORIDE AND POTASSIUM CHLORIDE.




FIGURE II.

same amount of potassium chloride is added to the succeeding mixtures, the calcium chloride can increase its water of hydration only slightly, and by a constant difference, with increase in the concentration of the calcium chloride added. Consequently, there is increasing deviation in Figure I between the curve representing the value of M for the single solution and that representing M for the mixture, that is, in Figure II, the difference between the values of M for calcium chloride in the separate solution and in the mixture is approximately proportional to the difference in the amount of water present acting as solvent toward the calcium chloride in the separate solution and in the mixture. This is shown by Curve II.

It must be borne in mind that these results are based on the assumption that calcium chloride breaks down at once into Ca^{++} , Cl^- , Cl^- . Also, that the entire decrease in conductivity upon mixing is due to re-formation of the molecules, whereas, a part of the decrease is certainly due to viscosity and friction of the ionic spheres, as already mentioned.

In Order to Ascertain the Effect upon the Hydrates Formed, a Pair of Salts, Both of Which Have Hydrating Power, Was Next Studied.

For this purpose it was considered desirable to in-

investigate, first, a case where both the salts employed have a large hydrating power, and where the hydrating power of the two salts in single solution is of the same order of magnitude. Calcium chloride and magnesium chloride fulfill these conditions.

CALCIUM CHLORIDE.

Table XVII. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ /m.
0.15	0.974	6.49
0.25	1.265	5.06
0.35	1.801	5.15
0.45	2.25	5.22
0.65	3.65	5.62
0.85	5.05	5.94
1.05	6.65	6.33
1.25	8.41	6.73
1.45	10.51	7.25

Table XVIII. CONDUCTIVITY MEASUREMENTS.

$$\Delta_{\infty}, 0^{\circ} = 147.5.$$

m.	v.	k.	Δ_{γ}	α .
0.15	6.6666	.014991	99.9	0.677
0.25	4.0000	.024077	96.8	.655
0.35	2.8571	.032710	93.4	.633
0.45	2.2222	.040688	88.2	.598
0.61	1.538	.051376	83.4	.565
0.85	1.1765	.066903	78.7	.544
1.05	0.9523	.078244	74.5	.505
1.25	0.8000	.088853	71.1	.482
1.45	0.6897	.097385	67.2	.455

Table XIX. WEIGHT-NORMAL CORRECTIONS.

m.	W _{Sol.}	W _{salt.}	W _{H₂O}	Correction Per Cent.
0.15	25.3227	0.11625	24.90645	0.4
0.25	25.5423	.09375	24.84855	0.6
0.35	25.6040	.07125	24.83275	0.7
0.45	26.0110	1.34875	24.76225	1.0
0.65	26.1364	1.80375	24.63465	1.5
0.85	26.8816	2.75875	24.52285	1.9
1.05	27.2846	2.91375	24.37085	2.5
1.25	27.6792	3.06875	24.21045	3.2
1.45	28.1293	4.02375	24.10555	3.6

Table XX.

HYDRATES.

m.	α .	L.	Δ/m .	L'.	H.	H.
0.15	0.677	4.58	4.93	4.91	6.0	40.2
0.25	.653	4.39	5.06	5.03	6.9	32.8
0.35	.633	4.32	5.15	5.11	9.7	27.3
0.45	.598	4.08	5.22	5.17	11.7	25.9
0.65	.565	3.96	5.62	5.53	15.7	24.3
0.85	.534	3.85	5.94	5.83	18.9	22.3
1.05	.505	3.74	6.33	6.18	21.9	20.9
1.25	.482	3.65	6.73	6.50	24.3	19.5
1.45	.455	3.55	7.25	6.99	27.3	18.8

MAGNESIUM CHLORIDE.Table XXI. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ/m .
0.15	0.747	4.98
0.25	1.306	5.22
0.75	1.910	5.46
0.45	2.537	5.64
0.65	3.854	5.93
0.85	5.404	6.36
1.05	7.181	6.84
1.25	9.236	7.39
1.45	11.47	7.91

TABLE 1

TABLE 2

α	β	γ	δ	ϵ	ζ	η
0.00	0.00	0.00	0.00	0.00	0.00	0.00
0.05	0.05	0.05	0.05	0.05	0.05	0.05
0.10	0.10	0.10	0.10	0.10	0.10	0.10
0.15	0.15	0.15	0.15	0.15	0.15	0.15
0.20	0.20	0.20	0.20	0.20	0.20	0.20
0.25	0.25	0.25	0.25	0.25	0.25	0.25
0.30	0.30	0.30	0.30	0.30	0.30	0.30
0.35	0.35	0.35	0.35	0.35	0.35	0.35
0.40	0.40	0.40	0.40	0.40	0.40	0.40
0.45	0.45	0.45	0.45	0.45	0.45	0.45
0.50	0.50	0.50	0.50	0.50	0.50	0.50

TABLE 3

TABLE 4

θ	ϕ	ψ
0.00	0.00	0.00
0.05	0.05	0.05
0.10	0.10	0.10
0.15	0.15	0.15
0.20	0.20	0.20
0.25	0.25	0.25
0.30	0.30	0.30
0.35	0.35	0.35
0.40	0.40	0.40
0.45	0.45	0.45
0.50	0.50	0.50

Table XXII. CONDUCTIVITY IN AQUEOUS SOLUTIONS.

m.	v.	P.	Λ_v .	α .
0.15	6.6666	.013643	90.9	0.705
0.25	4.0000	.021529	86.1	.667
0.35	2.8571	.0302	80.7	.635
0.45	2.2222	.036183	80.1	.622
0.65	1.5385	.047864	73.6	.570
0.85	1.1765	.057606	67.8	.525
1.05	0.9523	.066280	63.1	.486
1.25	0.8000	.073499	58.8	.450
1.45	0.6896	.078040	53.8	.417

Table XXIII. WEIGHT-NORMAL CORRECTIONS.

m.	WSol.	WSalt.	WH ₂ O	Correction Per Cent.
0.15	25.2154	0.7573	24.9332	0.3
0.25	25.5072	.5954	24.9118	.4
0.35	25.7066	.4375	24.8751	.5
0.45	25.8825	1.717	24.8108	.8
0.65	26.2378	1.5180	24.6698	1.2
0.85	26.6460	2.0245	24.6217	1.5
1.05	27.0011	2.5006	24.5005	2.0
1.25	27.3538	2.9768	24.3770	2.5
1.45	27.7095	3.4532	24.2563	3.0

TABLE 1. SUMMARY OF RESULTS

Case	Year	Age	Sex	Site
1	1950	20	M	1
2	1951	21	M	1
3	1952	22	M	1
4	1953	23	M	1
5	1954	24	M	1
6	1955	25	M	1
7	1956	26	M	1
8	1957	27	M	1
9	1958	28	M	1
10	1959	29	M	1
11	1960	30	M	1
12	1961	31	M	1

TABLE 2. SUMMARY OF RESULTS

Case	Year	Age	Sex	Site
1	1950	20	M	1
2	1951	21	M	1
3	1952	22	M	1
4	1953	23	M	1
5	1954	24	M	1
6	1955	25	M	1
7	1956	26	M	1
8	1957	27	M	1
9	1958	28	M	1
10	1959	29	M	1
11	1960	30	M	1
12	1961	31	M	1

Table XXIV.

HYDRATES.

m.	α .	L.	Δ/m .	L'.	M.	H.
0.15	0.705	4.46	4.96	4.97	5.5	26.5
0.25	.667	4.24	5.23	5.20	6.2	26.7
0.35	.625	4.19	5.46	5.43	12.7	26.7
0.45	.601	4.10	5.64	5.69	14.9	23.1
0.65	.570	3.98	5.93	5.86	17.8	27.4
0.85	.535	3.81	6.36	6.26	21.7	25.6
1.05	.489	3.68	6.84	6.70	25.1	23.9
1.25	.456	3.56	7.39	7.20	28.1	22.5
1.45	.417	3.41	7.91	7.67	30.9	21.3

MIXTURE OF CALCIUM CHLORIDE AND MAGNESIUM CHLORIDE.

Table XXV. FREEZING-POINT MEASUREMENTS.

m.	P.	Δ .
0.3 CaCl_2	1	1.553
0.5 MgCl_2	1	
0.5 CaCl_2	1	2.704
0.5 MgCl_2	"	
0.7 CaCl_2	1	4.053
0.7 MgCl_2	"	
0.9 CaCl_2	1	5.543
0.9 MgCl_2	"	
1.3 CaCl_2	1	9.320
1.3 MgCl_2	"	
1.7 CaCl_2	1	12.50
1.7 MgCl_2	"	
2.1 CaCl_2	1	16.50
2.1 MgCl_2	"	
2.5 CaCl_2	1	27.25
2.5 MgCl_2	"	
2.9 CaCl_2	1	37.5
2.9 MgCl_2	"	

TABLE 1 Summary of Data

Year	1950	1951	1952	1953	1954	1955
Population	100,000	105,000	110,000	115,000	120,000	125,000
Area	100	105	110	115	120	125
Population Density	1,000	1,048	1,091	1,130	1,167	1,200
Area Density	1.0	1.05	1.1	1.15	1.2	1.25
Population per Area	10,000	10,480	10,910	11,300	11,670	12,000
Area per Population	0.0001	0.000095	0.000091	0.000087	0.000083	0.00008
Population per Area Density	10,000	10,480	10,910	11,300	11,670	12,000
Area per Population Density	0.0001	0.000095	0.000091	0.000087	0.000083	0.00008
Population per Area Density	10,000	10,480	10,910	11,300	11,670	12,000
Area per Population Density	0.0001	0.000095	0.000091	0.000087	0.000083	0.00008

NOTE: The above data are based on the 1950 Census of the United States.

TABLE 2 Summary of Data

Year	1950	1951	1952	1953	1954	1955
Population	100,000	105,000	110,000	115,000	120,000	125,000
Area	100	105	110	115	120	125
Population Density	1,000	1,048	1,091	1,130	1,167	1,200
Area Density	1.0	1.05	1.1	1.15	1.2	1.25
Population per Area	10,000	10,480	10,910	11,300	11,670	12,000
Area per Population	0.0001	0.000095	0.000091	0.000087	0.000083	0.00008
Population per Area Density	10,000	10,480	10,910	11,300	11,670	12,000
Area per Population Density	0.0001	0.000095	0.000091	0.000087	0.000083	0.00008
Population per Area Density	10,000	10,480	10,910	11,300	11,670	12,000
Area per Population Density	0.0001	0.000095	0.000091	0.000087	0.000083	0.00008

NOTE: The above data are based on the 1950 Census of the United States.

TABLE XXVI. CONDUCTIVITY MEASUREMENTS.

μ	P	η_c	V_c	κ_w	κ	D	κ_c	Δ_c	α
0.8 CaCl_2	1	.15	3.6066	.028323	.014331		.014082	95.3	0.826
0.9 LiCl_2	1	.15	"		.013842	.001711		85.7	.861
0.5 CaCl_2	1	.25	4.000		.024773		.081703	96.9	.895
0.5 LiCl_2	1	.25	"	.041214	.021523	.004338	.013505	78.0	.805
0.7 CaCl_2	1	.35	2.8571	.053615	.022730	.007325	.023317	22.3	.553
0.7 LiCl_2	1	.35	"		.028240		.024738	70.3	.543
0.3 CaCl_2	1	.45	2.2222	.085030	.029633	.010731	.033322	75.2	.510
0.3 LiCl_2	1	.45	"		.026132		.031253	62.5	.532
1.3 CaCl_2	"	.65	1.333		.054236		.043253	62.3	.451
1.3 LiCl_2	1	.85	"	.032101	.047364	.013013	.033575	50.3	.430
1.7 CaCl_2	"	.85	1.1765	.034333	.066372	.033017	.050322	52.3	.402
1.7 LiCl_2	1	.25	"		.057606		.045022	51.1	.526
2.1 CaCl_2	"	1.05	0.9523		.072044		.055224	52.2	.357
2.1 LiCl_2	1	1.05	"	.10078	.063020	.04874	.04552	42.4	.532
3.5 CaCl_2	1	1.25	0.8000	.10273	.098853	.05357	.057313	46.3	.314
2.5 LiCl_2	"	1.25	"		.072430		.044332	35.3	.073
2.3 CaCl_2	1	1.45	0.63366		.037335		.053345	40.6	.275
2.3 LiCl_2	"	1.45	"	.10007	.072010	.07533	.041233	29.4	.220

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

Table 1.1. Summary of the data.

Table XVII. WEIGHT-NORMAL CORRECTIONS.

n.	P.	W _{sol.}	W _{salt.}	W _{H₂O}	Correction Per Cent.
0.5 CaCl ₂ 0.5 MgCl ₂	1 "	25.6460	0.77345	24.67255	0.5
0.5 CaCl ₂ 0.5 MgCl ₂	1 "	26.0814	1.23915	24.74225	1.0
0.7 CaCl ₂ 0.7 MgCl ₂	1 "	26.4594	1.80475	24.65465	1.4
0.9 CaCl ₂ 0.9 MgCl ₂	1 "	26.8453	2.32045	24.52485	1.9
1.3 CaCl ₂ 1.3 MgCl ₂	1 "	27.6088	3.35175	24.25705	3.0
1.7 CaCl ₂ 1.7 MgCl ₂	1 "	28.3321	4.38305	23.94905	4.0
2.1 CaCl ₂ 2.1 MgCl ₂	1 "	29.1096	5.41435	23.69525	5.2
2.5 CaCl ₂ 2.5 MgCl ₂	1 "	29.8377	6.44555	23.38775	6.5
2.9 CaCl ₂ 2.9 MgCl ₂	1 "	30.5280	7.47695	23.05105	7.2

In Table XVIII, n = normal concentration of each salt in the mixture. α = dissociation of each salt in the mixture, as given in Table XVI. mL = lowering of freezing-point, calculated from formula $[(1.86 \times 2\alpha) + 1.86]n$. The correction per cent is deduced on the assumption that the two salts form the same hydrates in the mixture that

they form in separate solutions. Upon this assumption the amount of water which each salt would eliminate from the sphere of action as solvent is calculated from the formula:— $1000 - \left[\frac{1}{L} \times 1000 \right]$. The values for magnesium chloride and calcium chloride are added together and to this is added also the difference between 1000 grams and the amount of water actually present in one liter as given by the weight-normal corrections. This, then, is given as a percentage correction in column headed Correction Per Cent. The value mL is divided by this per cent, and the result given under

Table XXVIII.

m.	P.	m ₀ .	α .	mL.	Correction Per Cent	Δ .
0.3 CaCl ₂	1	0.15	.636	0.634	78.9	0.803
0.3 MgCl ₂	"		.664	0.650		0.825
0.5 CaCl ₂	1	0.25	.589	1.012	68.0	1.488
0.5 MgCl ₂	"		.605	1.027		1.510
0.7 CaCl ₂	1	0.35	.558	1.378	58.8	2.345
0.7 MgCl ₂	"		.549	1.366		2.325
0.9 CaCl ₂	1	0.45	.510	1.690	51.2	3.298
0.9 MgCl ₂	"		.538	1.738		3.392
1.3 CaCl ₂	1	0.65	.454	2.36	38.5	5.998
1.3 MgCl ₂	"		.460	2.321		6.035
1.7 CaCl ₂	1	0.85	.406	2.865	25.8	11.13
1.7 MgCl ₂	"		.396	2.832		11.00
2.1 CaCl ₂	1	1.05	.357	3.346	14.7	22.25
2.1 MgCl ₂	"		.336	3.261		22.29

Evidently the sum of the values of Δ for calcium ch-

loride and Magnesium chloride, as thus found, should equal the lowering found experimentally for the mixture in question. These values are placed side by side, for comparison, in

Table XXIX.

m.	P.	m _C .	Δ (calculated)	Δ (Exp)	Diff.
0.3 CaCl ₂	1	0.15	1.626	1.553	0.073
0.3 MgCl ₂	"	"			
0.5 CaCl ₂	1	0.25	2.998	2.704	0.294
0.5 MgCl ₂	"	"			
0.7 CaCl ₂	1	0.35	4.670	4.053	0.617
0.7 MgCl ₂	"	"			
0.9 CaCl ₂	1	0.45	6.690	5.543	1.147
0.9 MgCl ₂	"	"			
1.3 CaCl ₂	1	0.65	12.033	9.320	2.713
1.3 MgCl ₂	"	"			
1.7 CaCl ₂	1	0.85	22.13	13.5	8.639
1.7 MgCl ₂	"	"			
2.1 CaCl ₂	1	1.05	45.14	19.5	25.64
2.1 MgCl ₂	"	"			

It is unnecessary to extend these calculations to the remaining more concentrated mixtures, since it is evident that the differences between the values calculated on the basis that the two salts form the same hydrates in separate solutions that they form in the mixtures, and the values

for Δ obtained experimentally, become increasingly large with increase in concentration. It is farther evident that the hypothesis is untenable from the fact that it would call for the elimination of 944 grams of water per liter of solution in the case of the 2.5 N mixture of the above salts, and for 1047 grams eliminated from the sphere of action as solvent in the case of the 2.9 N mixture.

At the moment of mixing the two solutions, each salt is prevented from forming the hydrate which it ordinarily forms in separate solution, by the amount of water which has been eliminated from the sphere of action as solvent in the formation of the hydrate of the other component of the mixture. Consequently, for the more dilute solutions at least, the hydrates formed by a salt in separate solution should be related to the hydrates formed by the same salt in the mixture at corresponding concentrations, as the amount of water present as solvent in the single solution is to the amount present as solvent in the mixture. That is, taking calcium chloride and magnesium chloride, 0.15 N solutions of each, referring to Tables XX and XXIV, if x represents the unknown value of N in the mixture; for calcium chloride, $0.0 : x :: 1000 : 857$. 0.0 is the value of N for 0.15 N

the first of these is the fact that the
 the second is the fact that the
 the third is the fact that the
 the fourth is the fact that the
 the fifth is the fact that the
 the sixth is the fact that the
 the seventh is the fact that the
 the eighth is the fact that the
 the ninth is the fact that the
 the tenth is the fact that the

the eleventh is the fact that the
 the twelfth is the fact that the
 the thirteenth is the fact that the
 the fourteenth is the fact that the
 the fifteenth is the fact that the
 the sixteenth is the fact that the
 the seventeenth is the fact that the
 the eighteenth is the fact that the
 the nineteenth is the fact that the
 the twentieth is the fact that the
 the twenty-first is the fact that the
 the twenty-second is the fact that the
 the twenty-third is the fact that the
 the twenty-fourth is the fact that the
 the twenty-fifth is the fact that the
 the twenty-sixth is the fact that the
 the twenty-seventh is the fact that the
 the twenty-eighth is the fact that the
 the twenty-ninth is the fact that the
 the thirtieth is the fact that the
 the thirty-first is the fact that the
 the thirty-second is the fact that the
 the thirty-third is the fact that the
 the thirty-fourth is the fact that the
 the thirty-fifth is the fact that the
 the thirty-sixth is the fact that the
 the thirty-seventh is the fact that the
 the thirty-eighth is the fact that the
 the thirty-ninth is the fact that the
 the fortieth is the fact that the
 the forty-first is the fact that the
 the forty-second is the fact that the
 the forty-third is the fact that the
 the forty-fourth is the fact that the
 the forty-fifth is the fact that the
 the forty-sixth is the fact that the
 the forty-seventh is the fact that the
 the forty-eighth is the fact that the
 the forty-ninth is the fact that the
 the fiftieth is the fact that the
 the fifty-first is the fact that the
 the fifty-second is the fact that the
 the fifty-third is the fact that the
 the fifty-fourth is the fact that the
 the fifty-fifth is the fact that the
 the fifty-sixth is the fact that the
 the fifty-seventh is the fact that the
 the fifty-eighth is the fact that the
 the fifty-ninth is the fact that the
 the sixtieth is the fact that the
 the sixty-first is the fact that the
 the sixty-second is the fact that the
 the sixty-third is the fact that the
 the sixty-fourth is the fact that the
 the sixty-fifth is the fact that the
 the sixty-sixth is the fact that the
 the sixty-seventh is the fact that the
 the sixty-eighth is the fact that the
 the sixty-ninth is the fact that the
 the seventieth is the fact that the
 the seventy-first is the fact that the
 the seventy-second is the fact that the
 the seventy-third is the fact that the
 the seventy-fourth is the fact that the
 the seventy-fifth is the fact that the
 the seventy-sixth is the fact that the
 the seventy-seventh is the fact that the
 the seventy-eighth is the fact that the
 the seventy-ninth is the fact that the
 the eightieth is the fact that the
 the eighty-first is the fact that the
 the eighty-second is the fact that the
 the eighty-third is the fact that the
 the eighty-fourth is the fact that the
 the eighty-fifth is the fact that the
 the eighty-sixth is the fact that the
 the eighty-seventh is the fact that the
 the eighty-eighth is the fact that the
 the eighty-ninth is the fact that the
 the ninetieth is the fact that the
 the ninety-first is the fact that the
 the ninety-second is the fact that the
 the ninety-third is the fact that the
 the ninety-fourth is the fact that the
 the ninety-fifth is the fact that the
 the ninety-sixth is the fact that the
 the ninety-seventh is the fact that the
 the ninety-eighth is the fact that the
 the ninety-ninth is the fact that the
 the hundredth is the fact that the

calcium chloride in separate solution, and this has been corrected to 1000 grams of solvent. The figure 897 is found by subtracting from 1000 the number of grams of water with which the magnesium chloride combines when it forms the hydrate indicated for the concentration in question in Table XXIV, plus the difference between the amount of water present in the solution and 1000 grams as given by the weight-normal correction.

Having thus obtained the values of M for the salts in the mixture, it is possible to solve for L' from the equation

$$M = \frac{1000 - \left[\frac{L}{L'} \times 1000 \right]}{18}, \quad \text{where } L \text{ represents the}$$

theoretical lowering found for the salt by calculating it from the value of α which has been determined for the mixture in question. This value of L' must then be corrected, in order to obtain the value as it would be observed, by dividing it by the proper per cent. This percentage correction is found by deducing the number of grams of water eliminated as water of hydration by the other salt in the mixture, basing the calculation upon the value of M worked out for the mixture. To this must be added the percentage correction which the weight-normal correction has given.

The sum of these values for L' for the two salts in the mixture should then be equal to Δ as found experimentally for the mixture, provided this hypothesis is correct. In the following table the values thus obtained are given.

M_0 = the value for M deduced from the proportion.

L'_0 = " " " L' calculated upon this value of M .

$L'_0{}^{Cor.}$ = " " " L' , with proper percentage correction.

Sum = the sum of these two values for $L'_0{}^{Cor.}$ multiplied by the normality of the solution.

Δ = the lowering found experimentally for the same mixture.

The other symbols employed have the usual significance.

The first of these is the fact that the

the second is the fact that the
the third is the fact that the
the fourth is the fact that the
the fifth is the fact that the
the sixth is the fact that the
the seventh is the fact that the
the eighth is the fact that the
the ninth is the fact that the
the tenth is the fact that the

the eleventh is the fact that the

the twelfth is the fact that the

the thirteenth is the fact that the

the fourteenth is the fact that the

the fifteenth is the fact that the

the sixteenth is the fact that the

the seventeenth is the fact that the

TABLE XXX.

λ .	ρ .	n_c .	M .	M_c .	L .	L_0 .	L_0^{con} .	Sum.	Δ .
0.2 CaCl_2	1	.15	6.0	4.4	4.224	4.632	5.15	1.57	1.552
0.3 MgCl_2	"	.15	5.5	4.9	4.331	4.748	5.29		
0.5 CaCl_2	1	.25	3.2	6.9	4.109	4.635	5.52	2.74	2.704
0.5 MgCl_2	"	.25	9.2	7.9	4.05	4.715	5.44		
0.7 CaCl_2	1	.35	9.7	7.5	3.936	4.547	5.71	3.97	4.059
0.7 MgCl_2	"	.35	13.7	10.5	3.902	4.808	5.64		
0.0 CaCl_2	1	.45	11.7	8.6	3.755	4.433	5.77	5.22	5.545
0.9 MgCl_2	"	.45	14.9	11.9	3.832	4.397	5.32		
1.2 CaCl_2	1	.65	15.7	10.7	3.548	4.394	5.94	7.74	9.22
1.2 MgCl_2	"	.65	17.3	12.3	3.57	4.633	5.97		
1.7 CaCl_2	1	.85	19.9	11.5	3.371	4.251	6.06	10.21	13.5
1.7 MgCl_2	"	.85	21.7	14.3	3.332	4.437	5.96		

一 二 三 四 五 六 七 八

九 十 十一 十二 十三 十四 十五 十六

十七 十八 十九 二十 二十一 二十二 二十三 二十四

二十五 二十六 二十七 二十八 二十九 三十 三十一 三十二

三十三 三十四 三十五 三十六 三十七 三十八 三十九 四十

四十一 四十二 四十三 四十四 四十五 四十六 四十七 四十八

四十九 五十 五十一 五十二 五十三 五十四 五十五 五十六

五十七 五十八 五十九 六十 六十一 六十二 六十三 六十四

六十五 六十六 六十七 六十八 六十九 七十 七十一 七十二

七十三 七十四 七十五 七十六 七十七 七十八 七十九 八十

Without further extending this method of calculation it is evident that it does not give the true value of the hydrates formed, and that the hydrating power is larger for the more concentrated mixtures than would be indicated by the proportions given above.

This is just what we should expect, for with increasing hydration we do not have the process of loss of water from the hydrates formed in separate solution going on to the end in the mixture, but instead we have, necessarily, a condition of equilibrium between the hydrating powers of the two salts. Also, as a direct result, the amount of water thus left uncombined affects the final composition of the hydrates, as already indicated.

This condition of equilibrium is not represented by the proportions given above, since the amount of water playing the part of solvent is not that indicated by these proportions.

If we consider that the values of M for calcium chloride and for magnesium chloride in separate solutions, since their hydrating powers are nearly equal, represent the relative hydrating powers under the same conditions for the two salts, at that concentration corresponding to

any pair of values of M , then it becomes possible to calculate the approximate composition of the hydrates of calcium chloride and magnesium chloride in the mixture, for, selecting for example the mixture consisting of 1 part of 1.7 N calcium chloride and 1 part of 1.7 N magnesium chloride, then:-

Let x equal the number of grams of water eliminated in formation of the hydrate of calcium chloride which occurs in the mixture in question. If the hydrates of calcium chloride and magnesium chloride in the mixture are related as they are in separate solutions, they will be to each other as 18.9 : 21.7 (see Tables XX and XXIV), hence,

$\frac{217}{189}x$ = the number of grams of water eliminated in formation of hydrate of magnesium chloride in the mixture.

From formula, $M = \frac{1000 - \left[\frac{L}{I} \times 1000 \right]}{18}$, we have,

$$(1) \quad x = 1000 - \frac{3371}{L_{Ca}}, \text{ where } 3.371 \text{ is the lowering found for}$$

0.85 N calcium chloride by calculation from the value of α given in Table XXVI, and L_{Ca}' = the lowering observed for calcium chloride corrected properly for water lacking from 1000 grams, and multiplied by m .

$$\text{Similarly, } \frac{217}{189}x = 1000 - \frac{5332}{L_{Mg}'}. \quad (2)$$

Let L'_0 = observed lowering due to magnesium chloride.

Then, $18.5 - L'_0$ = observed lowering due to calcium chloride, (vld. Table XXV).

$$\text{Then, } L'_0 = (L'_{Mg} \times 0.85) \div \left(1 - \frac{x}{1000} - \frac{4}{100}\right). \quad (3)$$

where 0.85 is the normality in question, and the second parenthesis is the necessary percentage correction, based upon the value of x and the correction given in Table XXVII.

$$\text{Similarly, } 18.5 - L'_0 = (L'_{Ca} \times 0.85) \div \left(1 - \frac{217x}{1000} - \frac{4}{100}\right). \quad (4)$$

$$\text{From equation (1), } L'_{Ca} = \frac{4371}{1000 - x}. \quad (5)$$

$$\text{" " (2), } L'_{Mg} = \frac{3372}{1.000 - 1.148x}. \quad (6)$$

Eliminating L'_0 by combining equations (3) and (4) and then substituting the values of L'_{Ca} and L'_{Mg} from (5) and (6), it becomes possible to solve for x . By this method the values of M for calcium chloride and magnesium chloride in this mixture are found to be 17.1 and 19.6, respectively.

By the above method the values of M for the mixtures between 0.9 N and 2.9 N have been derived, and are given in the following table, under L'_0 . The abbreviations at the top of each column have the usual significance, the suffix c indicating that the values are for the salt in the mixture.

Let f be a function defined on the interval $[a, b]$.

Then, f is continuous at a if and only if $\lim_{x \rightarrow a^+} f(x) = f(a)$.

$$\lim_{x \rightarrow a^+} f(x) = f(a) \iff \lim_{x \rightarrow a^+} f(x) = f(a) \text{ and } \lim_{x \rightarrow a^+} f(x) = f(a).$$

Now, let f be a function defined on the interval $[a, b]$. Then, f is continuous at a if and only if $\lim_{x \rightarrow a^+} f(x) = f(a)$.

Let f be a function defined on the interval $[a, b]$.

$$(a) \quad \lim_{x \rightarrow a^+} f(x) = f(a) \iff \lim_{x \rightarrow a^+} f(x) = f(a) \text{ and } \lim_{x \rightarrow a^+} f(x) = f(a).$$

$$(b) \quad \lim_{x \rightarrow a^+} f(x) = f(a) \iff \lim_{x \rightarrow a^+} f(x) = f(a) \text{ and } \lim_{x \rightarrow a^+} f(x) = f(a).$$

$$(c) \quad \lim_{x \rightarrow a^+} f(x) = f(a) \iff \lim_{x \rightarrow a^+} f(x) = f(a) \text{ and } \lim_{x \rightarrow a^+} f(x) = f(a).$$

Let f be a function defined on the interval $[a, b]$. Then, f is continuous at a if and only if $\lim_{x \rightarrow a^+} f(x) = f(a)$.

Let f be a function defined on the interval $[a, b]$. Then, f is continuous at a if and only if $\lim_{x \rightarrow a^+} f(x) = f(a)$.

Let f be a function defined on the interval $[a, b]$.

Let f be a function defined on the interval $[a, b]$. Then, f is continuous at a if and only if $\lim_{x \rightarrow a^+} f(x) = f(a)$.

Let f be a function defined on the interval $[a, b]$. Then, f is continuous at a if and only if $\lim_{x \rightarrow a^+} f(x) = f(a)$.

Let f be a function defined on the interval $[a, b]$. Then, f is continuous at a if and only if $\lim_{x \rightarrow a^+} f(x) = f(a)$.

TABLE XXXI.

L.	F.	ϵ_e .	α .	ϵ_e .	λ .	Δ .	$\epsilon_{\text{Gr.}}$.	λ_e .	μ_e .
1.5	CaCl_2 1	0.65	0.454	3.549	15.74	9° 32	3.0	14.5	22.5
1.5	MgCl_2 "	0.65	.460	3.57	17.79			16.4	25.2
1.7	CaCl_2 1	0.35	.406	3.371	13.92	15.5	4.0	17.1	20.1
1.7	MgCl_2 "	0.35	.396	3.332	21.73			19.6	25.1
2.1	CaCl_2 1	1.05	.357	3.137	21.92	19.5	5.2	19.3	16.9
2.1	MgCl_2 "	1.05	.336	3.264	25.06			22.7	21.6
2.5	CaCl_2 1	1.25	.314	3.023	24.32	27.25	6.5	22.3	13.2
2.5	MgCl_2 "	1.25	.275	2.935	28.14			26.9	21.0
2.9	CaCl_2 1	1.45	.275	2.383	27.29	37.0	7.3	25.9	17.9
2.9	MgCl_2 "	1.45	.220	2.679	30.86			28.9	19.9

" " MIXED WITH CALCIUM CHLORIDE.
 " " CALCIUM CHLORIDE ALONE.
 " " MIXED WITH MAGNESIUM CHLORIDE.

TOTAL NUMBER OF GRAM-MOLECULES OF WATER COMPARED.




FIGURE III.

For the sake of comparison the values of M are plotted as curves for calcium chloride and magnesium chloride in separate solutions against concentration, (as axis of abscissas). Upon the same sheet are drawn similar curves for the values of M in the mixture.

Upon comparing curve I, the hydration of magnesium chloride alone, with curve IV, the hydration of the calcium chloride in the mixture, it will be seen that where I rises IV falls, and vice versa. This is exactly what we should expect, since an increase in the hydrating power of magnesium chloride should result in a corresponding decrease in the amount of water with which the calcium chloride would be able to combine in the mixture, unless a corresponding increase in the hydrating power of the calcium chloride in separate solution should also occur. Again, comparing curves I and II, and III and IV, the hydrates in separate solution and in the mixture are of the same general order of magnitude, except where the presence of the one salt in the mixture alters the composition of the hydrate formed by the other.

It was Thought Desirable to Compare the Effect of a Salt with less Hydrating Power Mixed with One of Greater Hydrating Power.

For this purpose calcium chloride and strontium chlor

The last part of the paper is devoted to the study of the asymptotic behavior of the solutions of the system (1) as $t \rightarrow \infty$. It is shown that the solutions of the system (1) are bounded and tend to zero as $t \rightarrow \infty$. The proof is based on the use of the Lyapunov function method.

Let us consider the system (1) in the case when $\lambda = 0$. In this case the system (1) can be written in the form

$$\dot{x} = -x, \quad \dot{y} = -y, \quad \dot{z} = -z, \quad \dot{w} = -w, \quad \dot{v} = -v, \quad \dot{u} = -u, \quad \dot{t} = 1.$$

The solutions of this system are

$$x(t) = x_0 e^{-t}, \quad y(t) = y_0 e^{-t}, \quad z(t) = z_0 e^{-t}, \quad w(t) = w_0 e^{-t}, \quad v(t) = v_0 e^{-t}, \quad u(t) = u_0 e^{-t}, \quad t(t) = t_0 + t.$$

It is easy to see that the solutions of the system (1) are bounded and tend to zero as $t \rightarrow \infty$. The proof is based on the use of the Lyapunov function method.

It is easy to see that the solutions of the system (1) are bounded and tend to zero as $t \rightarrow \infty$. The proof is based on the use of the Lyapunov function method.

ide were selected, the hydrating power of strontium chloride being somewhat less than that of calcium chloride.

The data for the separate solutions of calcium chloride has already been given. That of strontium chloride follows.

STRONTIUM CHLORIDE.

Table XXXII. FREEZING-POINT MEASUREMENTS.

n.	Δ .	Δ/m .
0.15	0.753	5.02
0.25	1.249	5.00
0.35	1.774	5.07
0.45	2.233	5.18
0.65	3.521	5.42
0.85	4.863	5.72
1.05	6.354	6.07
1.25541	7.861	6.56

Table XXXIII. CONDUCTIVITY MEASUREMENTS.

m.	v.	k.	Λ_v .	α .
0.15	0.6666	.015377	102.5	.747
0.25	1.0000	.024185	96.7	.705
0.35	2.8571	.02259	93.1	.678
0.45	2.2222	.010479	90.0	.655
0.65	1.538	.055017	84.6	.616
0.85	1.1764	.067824	79.8	.581
1.05	0.9523	.079796	76.0	.553
1.25541	0.8945	.08607	71.7	.522

For more details, see the Appendix, Part 2 of this report.

Table 1 shows the results of the analysis.

The data for the analysis are given in the following.

The results of the analysis are given in the following.

Table 1. Results of the analysis.

Table 1. Results of the analysis.		
Year	Month	Day
1970	1	1
1970	2	1
1970	3	1
1970	4	1
1970	5	1
1970	6	1
1970	7	1
1970	8	1
1970	9	1
1970	10	1
1970	11	1
1970	12	1

Table 2. Results of the analysis.

Year	Month	Day	Time	Location
1970	1	1	10:00	10:00
1970	2	1	10:00	10:00
1970	3	1	10:00	10:00
1970	4	1	10:00	10:00
1970	5	1	10:00	10:00
1970	6	1	10:00	10:00
1970	7	1	10:00	10:00
1970	8	1	10:00	10:00
1970	9	1	10:00	10:00
1970	10	1	10:00	10:00
1970	11	1	10:00	10:00
1970	12	1	10:00	10:00

Table XXXIV. WEIGHT-CORRECTED COMPOSITIONS.

m.	W _{sol.}	W _{salt.}	W _{H₂O.}	Correction Per Cent.
0.15	25.5325	0.59437	31.9401	0.8
0.35	25.8496	0.9906	24.85898	0.6
0.35	26.1900	1.52628	24.8071	0.8
0.45	26.5383	1.7871	24.7557	1.0
0.65	27.1944	3.5756	24.6128	1.5
0.85	27.8553	3.3681	24.4872	2.1
1.05	28.5672	4.1606	24.4066	2.4
1.23541	29.1297	4.3954	24.2344	3.1

Table XXXV. HYDRATES.

m.	α .	L.	Δ/m .	L'.	M.	H.
0.15	0.747	4.64	5.02	4.99	3.9	26.0
0.25	.705	4.48	5.00	4.97	5.5	22.0
0.35	.672	4.39	5.07	5.05	7.1	20.3
0.45	.655	4.30	5.18	5.13	9.0	20.0
0.65	.616	4.15	5.42	5.34	12.4	19.1
0.85	.581	4.02	5.72	5.60	15.7	18.5
1.05	.553	3.92	6.05	5.91	18.7	17.8
1.23541	.522	3.80	6.36	6.17	21.3	17.2

TABLE 1. SUMMARY OF DATA FOR THE 1960-1961 SEASON

STATION NO.	DATE	TIME	WIND	TEMP.	REL. HUM.
1.1	1960.1.15	10.00	10.0	10.0	10.0
1.2	1960.1.20	10.00	10.0	10.0	10.0
1.3	1960.1.25	10.00	10.0	10.0	10.0
1.4	1960.1.30	10.00	10.0	10.0	10.0
1.5	1960.2.5	10.00	10.0	10.0	10.0
1.6	1960.2.10	10.00	10.0	10.0	10.0
1.7	1960.2.15	10.00	10.0	10.0	10.0
1.8	1960.2.20	10.00	10.0	10.0	10.0
1.9	1960.2.25	10.00	10.0	10.0	10.0
1.10	1960.3.1	10.00	10.0	10.0	10.0

TABLE 2. SUMMARY OF DATA FOR THE 1961-1962 SEASON

STATION NO.	DATE	TIME	WIND	TEMP.	REL. HUM.
1.1	1961.1.15	10.00	10.0	10.0	10.0
1.2	1961.1.20	10.00	10.0	10.0	10.0
1.3	1961.1.25	10.00	10.0	10.0	10.0
1.4	1961.1.30	10.00	10.0	10.0	10.0
1.5	1961.2.5	10.00	10.0	10.0	10.0
1.6	1961.2.10	10.00	10.0	10.0	10.0
1.7	1961.2.15	10.00	10.0	10.0	10.0
1.8	1961.2.20	10.00	10.0	10.0	10.0
1.9	1961.2.25	10.00	10.0	10.0	10.0
1.10	1961.3.1	10.00	10.0	10.0	10.0

MIXTURE OF CALCIUM CHLORIDE AND STRONTIUM CHLORIDE.

Table XXXVI. FREEZING-POINT MEASUREMENTS.

n.	P.	Δ .
0.3 CaCl_2	1	1950°
0.3 SrCl_2	"	
0.5 CaCl_2	1	2.626
0.5 SrCl_2	"	
0.7 CaCl_2	1	3.906
0.7 SrCl_2	"	
0.9 CaCl_2	1	5.710
0.9 SrCl_2	"	
1.3 CaCl_2	1	8.682
1.3 SrCl_2	"	
1.7 CaCl_2	1	12.924
1.7 SrCl_2	"	
2.1 CaCl_2	1	18.29
2.1 SrCl_2	"	
2.5 CaCl_2	1	Salt separates out.
2.5 SrCl_2	"	

Table XXXVIII. WEIGHT-NORMAL CORRECTIONS.

n.	P.	$W_{\text{Sol.}}$	$W_{\text{Salt.}}$	$W_{\text{H}_2\text{O.}}$	Correction Per Cent.
0.3 CaCl_2	1	25.8463	1.0106	24.8357	0.7
0.3 SrCl_2	"				
0.5 CaCl_2	1	26.4106	1.6844	24.7262	1.1
0.5 SrCl_2	"				
0.7 CaCl_2	1	26.9452	2.3581	24.5871	1.7
0.7 SrCl_2	"				
0.9 CaCl_2	1	27.4711	3.0319	24.4392	2.2
0.9 SrCl_2	"				
1.3 CaCl_2	1	28.5671	4.7764	24.1877	3.5
1.3 SrCl_2	"				

TABLE 1. Summary of the results of the analysis of variance for the effect of the treatment on the response of the subjects.

TREATMENT		ANALYSIS OF VARIANCE		RESULTS	
Factor	Level	Source	SS	df	F
TREATMENT	1	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	2	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	3	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	4	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	5	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	6	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	7	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	8	Between	1.00	1	1.00
		Within	1.00	1	1.00

TABLE 2. Summary of the results of the analysis of variance for the effect of the treatment on the response of the subjects.

TREATMENT		ANALYSIS OF VARIANCE		RESULTS	
Factor	Level	Source	SS	df	F
TREATMENT	1	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	2	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	3	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	4	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	5	Between	1.00	1	1.00
		Within	1.00	1	1.00
TREATMENT	6	Between	1.00	1	1.00
		Within	1.00	1	1.00

TABLE XXXVII. CONDUCTIVITY MEASUREMENTS.

T.	P.	m.e.	V _c .	K _w .	K.	L.	K _c .	A _g .	α.
0.3 CoCl ₂	1	0.15	6.6666	.014931	.023239	.002109	.019917	32.5	.685
0.3 SrCl ₂	"	.15	"	.015877			.014400	36.0	.633
0.5 Ca	1	.25	4.0000	.024072	.043330	.004273	.021775	37.1	.530
0.5 Sr	"	.25	"	.024195			.022275	38.2	.547
0.7 Ca	1	.25	2.8571	.03270	.05188	.007102	.02330	39.5	.500
0.7 Sr	"	.25	"	.02253			.02202	38.7	.510
0.9 Ca	1	.45	2.2222	.039693	.070491	.003676	.034452	72.6	.513
0.9 Sr	"	.45	"	.040473			.028035	30.1	.520
1.3 Ca	1	.65	1.538	.054926	.037731	.013522	.044245	63.1	.491
1.3 Sr	"	.65	"	.055017			.046496	71.5	.501
1.7 Ca	1	.35	1.1765	.068903	.10555	.023177	.051170	81.9	.403
1.7 Sr	"	.95	"	.057824			.054900	134.0	.420
2.1 Ca	1	1.05	0.9523	.073244	.11429	.04876	.054534	59.0	.552
2.1 Sr	"	1.05	"	.073736			.053930	52.0	.414

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000	1001	1002	1003	1004	1005	1006	1007	1008	1009	1010	1011	1012	1013	1014	1015	1016	1017	1018	1019	1020	1021	1022	1023	1024	1025	1026	1027	1028	1029	1030	1031	1032	1033	1034	1035	1036	1037	1038	1039	1040	1041	1042	1043	1044	1045	1046	1047	1048	1049	1050	1051	1052	1053	1054	1055	1056	1057	1058	1059	1060	1061	1062	1063	1064	1065	1066	1067	1068	1069	1070	1071	1072	1073	1074	1075	1076	1077	1078	1079	1080	1081	1082	1083	1084	1085	1086	1087	1088	1089	1090	1091	1092	1093	1094	1095	1096	1097	1098	1099	1100	1101	1102	1103	1104	1105	1106	1107	1108	1109	1110	1111	1112	1113	1114	1115	1116	1117	1118	1119	1120	1121	1122	1123	1124	1125	1126	1127	1128	1129	1130	1131	1132	1133	1134	1135	1136	1137	1138	1139	1140	1141	1142	1143	1144	1145	1146	1147	1148	1149	1150	1151	1152	1153	1154	1155	1156	1157	1158	1159	1160	1161	1162	1163	1164	1165	1166	1167	1168	1169	1170	1171	1172	1173	1174	1175	1176	1177	1178	1179	1180	1181	1182	1183	1184	1185	1186	1187	1188	1189	1190	1191	1192	1193	1194	1195	1196	1197	1198	1199	1200	1201	1202	1203	1204	1205	1206	1207	1208	1209	1210	1211	1212	1213	1214	1215	1216	1217	1218	1219	1220	1221	1222	1223	1224	1225	1226	1227	1228	1229	1230	1231	1232	1233	1234	1235	1236	1237	1238	1239	1240	1241	1242	1243	1244	1245	1246	1247	1248	1249	1250	1251	1252	1253	1254	1255	1256	1257	1258	1259	1260	1261	1262	1263	1264	1265	1266	1267	1268	1269	1270	1271	1272	1273	1274	1275	1276	1277	1278	1279	1280	1281	1282	1283	1284	1285	1286	1287	1288	1289	1290	1291	1292	1293	1294	1295	1296	1297	1298	1299	1300	1301	1302	1303	1304	1305	1306	1307	1308	1309	1310	1311	1312	1313	1314	1315	1316	1317	1318	1319	1320	1321	1322	1323	1324	1325	1326	1327	1328	1329	1330	1331	1332	1333	1334	1335	1336	1337	1338	1339	1340	1341	1342	1343	1344	1345	1346	1347	1348	1349	1350	1351	1352	1353	1354	1355	1356	1357	1358	1359	1360	1361	1362	1363	1364	1365	1366	1367	1368	1369	1370	1371	1372	1373	1374	1375	1376	1377	1378	1379	1380	1381	1382	1383	1384	1385	1386	1387	1388	1389	1390	1391	1392	1393	1394	1395	1396	1397	1398	1399	1400	1401	1402	1403	1404	1405	1406	1407	1408	1409	1410	1411	1412	1413	1414	1415	1416	1417	1418	1419	1420	1421	1422	1423	1424	1425	1426	1427	1428	1429	1430	1431	1432	1433	1434	1435	1436	1437	1438	1439	1440	1441	1442	1443	1444	1445	1446	1447	1448	1449	1450	1451	1452	1453	1454	1455	1456	1457	1458	1459	1460	1461	1462	1463	1464	1465	1466	1467	1468	1469	1470	1471	1472	1473	1474	1475	1476	1477	1478	1479	1480	1481	1482	1483	1484	1485	1486	1487	1488	1489	1490	
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--

Table XXVIII (Cont.). WEIGHT-NORMAL CORRECTIONS

m.	P.	W _{Sol.}	W _{Salt.}	W _{H₂O.}	Correction Per Cent.	
1.7	CaCl ₂	1	29.6881	5.7869	2.9112	4.4
1.7	SrCl ₂	"				
3.1	CaCl ₂	1	30.6741	7.0744	33.6081	5.6
3.1	SrCl ₂	"				

For the sake of comparison, the lowerings of the freezing-point which would be obtained for the mixture of calcium chloride and strontium chloride are given in Table XXIX, basing them upon the same method of calculation as that already given in detail for Table XXI. The symbols used have the same significance in the two tables.

It will be noted that in Table XXIX, again, as in the case of the mixture of calcium chloride and magnesium chloride, the difference between the lowering as calculated upon the hypothesis that for the more dilute solutions at least, the hydrates formed by a salt in separate solution are related to the hydrates formed by the same salt in the mixture at corresponding concentrations, as the amount of water present as solvent in the mixture—the difference between this and the lowering found experimentally, becomes increasingly large. This has already been discussed in the preceding case.

TABLE XXXIX.

μ .	P.	μ_c .	M.	μ_c .	L.	L' .	L'_{cor} .	Sum.	Δ .
0.3 Ca	1	0.15	6.0	5.6	4.136	4.65	5.00	1.543	1.500
0.5 Sr	"	"	3.9	3.5	4.431	4.76	5.32		
0.5 Ca	1	0.25	3.2	7.3	4.053	4.67	5.16	2.646	2.322
0.5 Sr	"	"	5.5	4.6	4.266	4.65	5.43		
0.7 Ca	1	0.21	3.7	8.3	3.942	4.63	5.26	3.771	3.906
0.7 Sr	"	"	7.1	5.7	4.127	4.60	5.51		
0.9 Ca	1	0.45	11.7	3.5	3.791	4.57	5.36	4.973	6.370
0.9 Sr	"	"	3.0	6.9	4.03	4.50	5.70		
1.3 Ca	1	0.65	15.7	11.7	3.576	4.53	5.56	7.465	8.682
1.3 Sr	"	"	12.4	3.5	3.797	4.43	5.32		
7.7 Ca	1	0.35	13.0	12.3	3.373	4.39	5.61	3.877	12.924
7.7 Sr	"	"	15.7	3.7	3.593	4.25	6.00		
2.1 Ca	1	1.05	21.9	13.3	3.171	4.17	5.49	11.22	13.29
5.1 Sr	"	"	18.7	10.3	3.40	4.17	5.32		

The method of calculating the composition of the hydrates existing in a mixture of two salts with about the same hydrating power, such as calcium chloride and magnesium chloride, was based upon the assumption that the values of M in their separate solutions represent the relative hydrating powers under the same conditions for the two salts in their mixtures. While this applies where the hydrates formed are of approximately the same degree of complexity, it would involve an error in the case of two salts of very differing hydrating powers, for, if the power of forming hydrates, i.e. of eliminating water from the sphere of action as solvent, differs; then, when two such substances are mixed, if a decrease in the amount of water eliminated from the sphere of action as solvent occurs, this decrease must be assigned to the two salts inversely as their values of M in separate solutions. This follows from a consideration of the fact that the salt with greater hydrating power will have, consequently, greater power of dehydrating the salt with lesser hydrating power. Therefore, the salt which has less power to unite with water must suffer the greater loss in water of hydration in the presence of the salt with stronger hydrating power.

Upon this assumption the following method of calculating the approximate composition of the hydrates in such a

mixture, has been worked out. Take the case of the mixture consisting of one volume of 0.3 N strontium chloride and one volume of 0.3 N calcium chloride:-

Let x equal the total decrease in the number of grams of water eliminated from the sphere of action as solvent, in the mixture, as compared with that eliminated in the separate solutions.

Then $\frac{76}{99} x$ = the diminution in the number of grams of water eliminated by the calcium chloride, and

$\frac{60}{99} x$ = the diminution in the number of grams of water eliminated by the strontium chloride (see Tables XX and XXXV for the values of M in separate solutions).

108.5 grams of water were eliminated by the calcium chloride as water of hydration in the 0.15 N separate solution of that salt. (See Table XX). 70.5 grams were eliminated by the strontium chloride. (See Table XXXV). Therefore,

$$108.5 - 0.39x = 1000 - \frac{4186}{L_{Ca}^1}, \quad (1) \quad \text{and}$$

$$70.5 - 0.61x = 1000 - \frac{4461}{L_{Sr}^1}. \quad (2)$$

$$L_{Ca}^1 = \frac{4186}{891.5 + 0.39x}, \quad \text{and} \quad L_{Sr}^1 = \frac{4461}{929.5 + 0.61x}.$$

Let L_0^1 = observed lowering for calcium chloride in the mixture, then,

$$L_0^1 = \frac{4186 \times 0.15}{891.5 + 0.39x} + \frac{932.5 + 0.61x}{1000}, \quad (3) \quad \text{and}$$

$$1.503 - L_0^1 = \frac{4461 \times 0.15}{929.5 + 0.61x} - \frac{894.5 + 0.39x}{1000}. \quad (4)$$

$$\lim_{x \rightarrow 0} \frac{1}{x} = \infty, \quad \lim_{x \rightarrow 0} \frac{1}{x^2} = \infty.$$

Combining equations (3) and (4),

$$1.50 = \frac{4186 \times 0.15}{822 + 0.901x + .0002379x^2} + \frac{4461 \times 0.15}{822 + 0.902x + 0.002379x^2}.$$

From this equation the approximate value of x can be obtained.

The values of N and H for the mixture of calcium chloride and strontium chloride, thus obtained, are given in Table XL.

The symbols have the usual significance.

1874 Jan. 10. (1874) (1874) (1874)

1874 Jan. 10. (1874) (1874) (1874)

1874 Jan. 10. (1874) (1874) (1874)

1874 Jan. 10. (1874) (1874) (1874)

1874 Jan. 10. (1874) (1874) (1874)

1874 Jan. 10. (1874) (1874) (1874)

		Table		XL.		Δ	Sr. Gr. cor.	Ne.	He.
m.	P.	α .	Lo.	$\frac{M}{V}$.	$\frac{M}{V}$.				
0.5 CaCl ₂	1	0.625	4.186	6.0	6.0	1.503	0.7	5.1	34.0
0.5 SrCl ₂	"	.699	4.461	7.9	7.9			3.4	16.0
0.5 Ca	1	.590	4.056	8.2	8.2	2.026	1.1	7.3	29.2
0.5 Sr	"	.647	4.266	5.5	5.5			4.2	16.8
0.7 Ca	1	.560	3.942	9.7	9.7	3.906	1.6	9.3	26.1
0.7 Sr	"	.610	4.127	7.1	7.1			6.5	18.6
0.9 Ca	1	.519	3.791	11.7	11.7	5.310	2.2	11.1	24.7
0.9 Sr	"	.583	4.03	9.0	9.0			8.3	18.2
1.3 Ca	1	.461	3.576	15.7	15.7	8.682	3.3	15.1	23.2
1.3 Sr	"	.521	3.797	12.4	12.4			11.6	17.8
2.7 Ca	1	.408	3.378	18.9	18.9	12.924	4.4	18.3	21.5
2.7 Sr	"	.466	3.593	15.7	15.7			15.0	17.6
3.1 Ca	1	.352	3.171	21.9	21.9	18.39	5.6	21.5	20.5
3.1 Sr	"	.414	3.40	18.7	18.7			18.2	17.3

20 25 30 35 40 45 50

55 60 65 70 75 80 85

90 95 100 105 110 115 120

125 130 135 140 145 150 155

155 160 165 170 175 180 185

190 195 200 205 210 215 220

225 230 235 240 245 250 255

255 260 265 270 275 280 285

290 295 300 305 310 315 320

325 330 335 340 345 350 355

355 360 365 370 375 380 385

390 395 400 405 410 415 420

For the sake of comparison, the values of η for the separate solutions of calcium chloride and strontium chloride, and those in the mixture of the two salts, have been plotted in curves in Figure IV.

We find that the results expressed in Figure IV are of the same character as those shown in Figure III, for the mixture of calcium chloride and magnesium chloride. The curve of hydration of each salt in the mixture is parallel to the curve of its hydration in the single solution, except where large values as shown in Curve I, for calcium chloride alone, have produced a corresponding lessening in the amount of water which the strontium chloride has been able to unite with, in the mixture, at a normality of 0.15 to 0.35. An increase in the hydrating power of the calcium chloride must necessarily be accompanied by a corresponding decrease in the complexity of the hydrates of strontium chloride.


The results of the present study, which are in line with the findings of other studies, suggest that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study.

The first of the results of the present study is that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study. The second of the results of the present study is that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study. The third of the results of the present study is that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study. The fourth of the results of the present study is that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study. The fifth of the results of the present study is that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study. The sixth of the results of the present study is that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study. The seventh of the results of the present study is that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study. The eighth of the results of the present study is that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study. The ninth of the results of the present study is that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study. The tenth of the results of the present study is that the use of the present study is not only a valid method for the study of the present study, but also a valid method for the study of the present study.

CALCIUM CHLORIDE ALONE.
 " " MIXED WITH STRONTIUM CHLORIDE.
 STRONTIUM CHLORIDE ALONE.
 " " MIXED WITH CALCIUM CHLORIDE.

DEPT. EXPERIMENTAL ENGINEERING, BIBLEY COLLEGE, CORNELL UNIVERSITY.

TOTAL NUMBER OF GRAM MOLECULES OF WATER COMBINED.



B. G. CARPENTER, STRAUB N. 24

GRAM-MOLECULES OF SALT PER LITER OF SOLUTION.

FIGURE IV.

In order to compare the difference in the amount of water present as solvent in the single solution, and in the mixture with the difference in the value of M in the two cases; these values are tabulated. D_{H_2O} is the difference in the amount of water present as solvent in the separate solution and in the mixture. It is found for a given salt by adding to the number of grams of water eliminated as water of hydration by the other salt in the mixture, the difference between the correction to weight-normal standard in the case of the separate solution of the salt in question, and in the mixture. D_M is the difference between the value of M in the single solution and in the mixture.

Table XLI. CALCIUM CHLORIDE IN THE MIXTURE OF CALCIUM CHLORIDE AND STRONTIUM CHLORIDE.

m .	D_{H_2O} .	D_M .
0.15	46	0.9
0.25	81	0.9
0.35	127	0.4
0.45	162	0.6
0.65	226	0.6
0.85	295	0.6
1.05	359	0.4

Table XLII. STRONTIUM CHLORIDE IN MIXTURE OF CALCIUM CHLORIDE AND STRONTIUM CHLORIDE.

n .	D_{H_2O} .	D_m .
0.15	96	1.5
0.25	137	1.3
0.35	176	0.6
0.45	212	0.7
0.65	289	0.8
0.85	353	0.7
1.05	419	5

These values are plotted in curves in Figures V and VI. The concentrations are the abscissas, and for curve I the grams of water are taken for ordinates. In curve II gram-molecules of water form the ordinates.

TABLE 1. SUMMARY OF DATA FOR THE FIRST TWO YEARS OF THE
 STUDY OF THE EFFECTS OF THE 1964-65 DROUGHT ON THE
 GROWTH OF THE COMMON BEAN IN THE SOUTHERN PLAINS

Year	Area	Yield
1964	100	10.0
1965	100	10.0
1966	100	10.0
1967	100	10.0
1968	100	10.0
1969	100	10.0
1970	100	10.0
1971	100	10.0


The data in Table 1 show that the yield of the common bean in the Southern Plains was relatively stable during the first two years of the study. The yield was 10.0 bushels per acre in 1964 and 1965, and 10.0 bushels per acre in 1966 and 1967. The yield was 10.0 bushels per acre in 1968 and 1969, and 10.0 bushels per acre in 1970 and 1971. The yield was 10.0 bushels per acre in 1972 and 1973, and 10.0 bushels per acre in 1974 and 1975.

AND IN MIXTURE OF CALCIUM CHLORIDE AND STRONTIUM CHLORIDE.
 DIFFERENCE BETWEEN THE VALUES OF γ FOR CALCIUM CHLORIDE IN SINGLE SOLUTION AND
 IN THE MIXTURE OF CALCIUM CHLORIDE AND STRONTIUM CHLORIDE.

DEPT. EXPERIMENTAL ENGINEERING, SIBLEY COLLEGE, CORNELL UNIVERSITY.

I. GRAMS OF WATER.


II. GRAM MOLECULES OF WATER.



GRAM MOLECULES OF SALT PER LITER OF SOLUTION.
 FIGURE 7.

DIFFERENCE BETWEEN THE VALUES OF κ FOR STRONTIUM CHLORIDE IN SINGLE SOLUTION OF STRONTIUM CHLORIDE AND IN MIXTURE OF CALCIUM CHLORIDE AND STRONTIUM CHLORIDE.
 DIFFERENCE BETWEEN THE VALUES OF κ FOR STRONTIUM CHLORIDE IN SINGLE SOLUTION AND IN THE MIXTURE OF CALCIUM CHLORIDE AND STRONTIUM CHLORIDE.

DEPT. EXPERIMENTAL ENGINEERING, SIBLEY COLLEGE, CORNELL UNIVERSITY.



GRAM VOLUMES OF SALT PER LITER OF SOLUTION.

FIGURE VI.

It is readily seen from a comparison of the curves that the decrease in the amount of water present as solvent in the mixture, is not accompanied by a corresponding decrease in the values of M . Here again, as in the case of the mixture of calcium chloride and potassium chloride, the difference in the values of M is nearly constant, because the concentration of both salts present in the mixture is increased at the same rate. Slight changes might be manifest if the ions and molecules have different hydrating powers, as seems very probable. Then a decrease in the number of ions and a correspondingly large increase in the number of molecules of dissolved substance present, such as necessarily occurs owing to suppression of the ionization in the mixture, would be accompanied by a change in the number of grams of water combined. That this plays a part in the change noted seems very probable.

For purposes of comparison these same values are given for a few concentrations in the case of calcium chloride and magnesium chloride. The phenomena are of the same kind.

Table XLIII. CALCIUM CHLORIDE IN CALCIUM CHLORIDE AND MAGNESIUM CHLORIDE MIXTURE.

m .	D_{H_2O} .	D_M .
0.65	310.2	1.2
0.85	373.8	1.8
1.05	435.0	2.1
1.25	507.0	1.5
1.45	562.2	1.4

Table XLIV. MAGNESIUM CHLORIDE IN CALCIUM CHLORIDE AND MAGNESIUM CHLORIDE MIXTURE.

m.	$D_{H_2O}^D$	D_m
0.05	279	1.4
0.85	333	2.1
1.05	389	2.4
1.25	450	1.8
1.45	514	2.0

A Pair OF Nitrates Was Next Studied.

The salts must be of such a nature as not to unite to form complexes when mixed, and must have hydrating power. Magnesium nitrate and strontium nitrate were selected. The data for these two salts follows.

MAGNESIUM NITRATE.

Table XLV. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ/m .
0.15	0.749	4.99
0.25	1.252	5.01
0.35	1.805	5.16
0.45	2.395	5.32
0.55	2.919	5.49
0.65	3.609	5.64
0.75	4.293	5.86

TABLE 1. SUMMARY OF DATA FOR THE STUDY

TABLE 1. SUMMARY OF DATA FOR THE STUDY

Year	Area	Value
1980	100	100.0
1981	100	100.0
1982	100	100.0
1983	100	100.0
1984	100	100.0
1985	100	100.0

Source: Data from the study.

The data were obtained from the study.

The data were obtained from the study.

The data were obtained from the study.

The data were obtained from the study.

The data were obtained from the study.

TABLE 2. SUMMARY OF DATA FOR THE STUDY

Year	Area	Value
1980	100	100.0
1981	100	100.0
1982	100	100.0
1983	100	100.0
1984	100	100.0
1985	100	100.0

Table XVI. CONDUCTIVITY MEASUREMENTS.

$$\Lambda_{\infty}, 0^{\circ}, = 128.5.$$

m.	v.	k.	Λ_v .	α .
0.15	6.6666	.018701	91.3	0.711
0.25	4.0000	.021563	86.3	.671
0.35	2.8571	.026613	81.3	.636
0.45	2.2222	.035123	73.1	.607
0.55	1.8182	.041506	75.5	.587
0.65	1.538	.046942	73.2	.562
0.75	1.3333	.052260	69.7	.542

Table XVII. WEIGHT-NORMAL CORRECTIONS.

m.	$W_{\text{Sol.}}$	$W_{\text{Salt.}}$	$W_{\text{H}_2\text{O.}}$	Correction Per Cent.
0.15	25.4306	0.5567	24.8738	0.5
0.25	25.688	0.9277	24.7603	1.0
0.35	25.9782	1.2989	24.6794	1.3
0.45	26.2975	1.6610	24.5376	1.8
0.55	26.4748	2.0411	24.4336	2.3
0.65	26.7605	2.4122	24.3484	2.6
0.75	27.0203	2.7835	24.2371	3.1

TABLE 1. SUMMARY OF DATA FOR THE 1960-1961 SEASON

WATER = 1, 2, 3, 4

WATER	1	2	3	4
1960	1.00	1.00	1.00	1.00
1961	1.00	1.00	1.00	1.00
1962	1.00	1.00	1.00	1.00
1963	1.00	1.00	1.00	1.00
1964	1.00	1.00	1.00	1.00
1965	1.00	1.00	1.00	1.00
1966	1.00	1.00	1.00	1.00
1967	1.00	1.00	1.00	1.00

TABLE 2. SUMMARY OF DATA FOR THE 1962-1963 SEASON

WATER	1	2	3	4
1962	1.00	1.00	1.00	1.00
1963	1.00	1.00	1.00	1.00
1964	1.00	1.00	1.00	1.00
1965	1.00	1.00	1.00	1.00
1966	1.00	1.00	1.00	1.00
1967	1.00	1.00	1.00	1.00
1968	1.00	1.00	1.00	1.00
1969	1.00	1.00	1.00	1.00

Table XLVIII.

HYDRATES.

m.	α .	L.	Δ/m .	L'.	L.	H.
0.15	0.711	4.50	4.99	4.97	8.8	35.1
0.25	.671	4.36	5.01	4.96	6.7	26.9
0.35	.636	4.23	5.16	5.09	9.4	26.8
0.45	.607	4.12	5.32	5.22	11.7	26.0
0.55	.587	4.05	5.49	5.36	13.6	24.7
0.65	.562	3.95	5.68	5.49	15.6	24.0
0.75	.542	3.88	5.86	5.68	17.6	23.5

STRONTIUM NITRATE.Table XLIX. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ/d .
0.15	0.674	4.49
0.25	1.085	4.34
0.35	1.516	4.33
0.45	1.903	4.23
0.55	2.286	4.16
0.65	2.681	4.12
0.75	3.060	4.08

Table L. CONDUCTIVITY MEASUREMENTS.

m.	$\Lambda_{\infty}, 0^{\circ} = 134.4.$		Λ_v .	α .
	v.	k.		
0.15	6.6666	.018164	87.8	0.653
0.25	4.0000	.019905	79.6	.592
0.35	2.8571	.025744	73.6	.548
0.45	2.2222	.030645	68.1	.507
0.55	1.8182	.035106	63.3	.475
0.65	1.538	.038976	59.9	.446
0.75	1.3333	.042076	56.1	.417

INPUT DATAINPUT DATA

λ	β	γ	δ	ϵ	ζ	η
1.00	0.1	0.1	0.1	0.1	0.1	0.1
1.00	0.2	0.2	0.2	0.2	0.2	0.2
1.00	0.3	0.3	0.3	0.3	0.3	0.3
1.00	0.4	0.4	0.4	0.4	0.4	0.4
1.00	0.5	0.5	0.5	0.5	0.5	0.5
1.00	0.6	0.6	0.6	0.6	0.6	0.6
1.00	0.7	0.7	0.7	0.7	0.7	0.7
1.00	0.8	0.8	0.8	0.8	0.8	0.8
1.00	0.9	0.9	0.9	0.9	0.9	0.9
1.00	1.0	1.0	1.0	1.0	1.0	1.0

INPUT DATAINPUT DATA

λ	β	γ	δ	ϵ	ζ	η
1.00	0.1	0.1	0.1	0.1	0.1	0.1
1.00	0.2	0.2	0.2	0.2	0.2	0.2
1.00	0.3	0.3	0.3	0.3	0.3	0.3
1.00	0.4	0.4	0.4	0.4	0.4	0.4
1.00	0.5	0.5	0.5	0.5	0.5	0.5
1.00	0.6	0.6	0.6	0.6	0.6	0.6
1.00	0.7	0.7	0.7	0.7	0.7	0.7
1.00	0.8	0.8	0.8	0.8	0.8	0.8
1.00	0.9	0.9	0.9	0.9	0.9	0.9
1.00	1.0	1.0	1.0	1.0	1.0	1.0

INPUT DATA

$$A = \frac{1}{2} \left(\frac{1}{\lambda} + \frac{1}{\beta} \right)$$

λ	β	γ	δ	ϵ	ζ	η
1.00	0.1	0.1	0.1	0.1	0.1	0.1
1.00	0.2	0.2	0.2	0.2	0.2	0.2
1.00	0.3	0.3	0.3	0.3	0.3	0.3
1.00	0.4	0.4	0.4	0.4	0.4	0.4
1.00	0.5	0.5	0.5	0.5	0.5	0.5
1.00	0.6	0.6	0.6	0.6	0.6	0.6
1.00	0.7	0.7	0.7	0.7	0.7	0.7
1.00	0.8	0.8	0.8	0.8	0.8	0.8
1.00	0.9	0.9	0.9	0.9	0.9	0.9
1.00	1.0	1.0	1.0	1.0	1.0	1.0

Table II. WEIGHT-NORMAL CORRECTIONS.

m.	W _{Sol.}	W _{Salt.}	W _{H₂O.}	Correction Per Cent.
0.15	25.6245	0.7138	24.8307	0.7
0.25	26.0440	1.3512	24.6848	1.2
0.35	26.4408	1.8960	24.5449	1.6
0.45	26.8625	2.4576	24.4047	2.5
0.55	27.389	2.9793	24.5097	2.8
0.65	27.9907	3.5210	24.1697	3.3
0.75	28.0218	4.0627	24.0191	3.9

Table LII. HYDRATES.

m.	α .	L.	Δ/m .	L'.	M.	H.
0.15	0.655	4.29	4.49	4.46	2.1	14.1
0.25	.592	4.06	4.34	4.29	2.92	11.9
0.35	.548	3.80	4.33	4.25	4.6	13.2
0.45	.507	3.75	4.23	4.13	5.1	11.1
0.55	.475	3.67	4.16	4.04	5.6	10.3
0.65	.446	3.52	4.12	3.98	6.4	9.9
0.75	.417	3.41	4.08	3.92	7.2	9.6

The data for the mixture of magnesium nitrate and strontium nitrate follows.

TABLE 1. SUMMARY OF DATA FOR THE 1960-1961 SEASON

STATION NO.	STATION NAME	STATION TYPE	STATION CLASS	STATION STATUS
1	1000	1000	1000	1000
2	2000	2000	2000	2000
3	3000	3000	3000	3000
4	4000	4000	4000	4000
5	5000	5000	5000	5000
6	6000	6000	6000	6000
7	7000	7000	7000	7000
8	8000	8000	8000	8000
9	9000	9000	9000	9000

TABLE 2. SUMMARY OF DATA FOR THE 1962-1963 SEASON

STATION NO.	STATION NAME	STATION TYPE	STATION CLASS	STATION STATUS
1	1000	1000	1000	1000
2	2000	2000	2000	2000
3	3000	3000	3000	3000
4	4000	4000	4000	4000
5	5000	5000	5000	5000
6	6000	6000	6000	6000
7	7000	7000	7000	7000
8	8000	8000	8000	8000
9	9000	9000	9000	9000

For more information on the data for the 1960-1961 season, see the report for the 1960-1961 season.

For more information on the data for the 1962-1963 season, see the report for the 1962-1963 season.

Table LIII. FUSING-POINT MEASUREMENTS.

m.	P.	Δ .
0.3 $\text{Mg}(\text{NO}_3)_2$	1	1.9304
0.3 $\text{Sr}(\text{NO}_3)_2$	"	
0.5 $\text{Mg}(\text{NO}_3)_2$	1	2.751
0.5 $\text{Sr}(\text{NO}_3)_2$	"	
0.7 $\text{Mg}(\text{NO}_3)_2$	1	3.571
0.7 $\text{Sr}(\text{NO}_3)_2$	"	
0.9 $\text{Mg}(\text{NO}_3)_2$	1	4.573
0.9 $\text{Sr}(\text{NO}_3)_2$	"	
1.1 $\text{Mg}(\text{NO}_3)_2$	1	5.646
1.1 $\text{Sr}(\text{NO}_3)_2$	"	
1.3 $\text{Mg}(\text{NO}_3)_2$	1	6.893
1.3 $\text{Sr}(\text{NO}_3)_2$	"	
1.5 $\text{Mg}(\text{NO}_3)_2$	1	8.109
1.5 $\text{Sr}(\text{NO}_3)_2$	"	

Table LIV. WEIGHT-NORMAL CORRECTIONS.

m.	P.	W Sol.	W Salt.	W H_2O .	Correction Per Cent.
0.3 $\text{Mg}(\text{NO}_3)_2$	1				
0.3 $\text{Sr}(\text{NO}_3)_2$	"	26.044	1.8505	24.6926	1.2
0.5 $\text{Mg}(\text{NO}_3)_2$	1				
0.5 $\text{Sr}(\text{NO}_3)_2$	"	26.7087	2.3819	24.4268	2.3
0.7 $\text{Mg}(\text{NO}_3)_2$	1				
0.7 $\text{Sr}(\text{NO}_3)_2$	"	27.7970	3.1948	24.3022	3.2
0.9 $\text{Mg}(\text{NO}_3)_2$	1				
0.9 $\text{Sr}(\text{NO}_3)_2$	"	28.0570	4.1076	24.9494	4.2
1.1 $\text{Mg}(\text{NO}_3)_2$	1				
1.1 $\text{Sr}(\text{NO}_3)_2$	"	28.6884	5.0308	24.6681	5.7
1.3 $\text{Mg}(\text{NO}_3)_2$	1				
1.3 $\text{Sr}(\text{NO}_3)_2$	"	29.7503	5.9732	24.4177	6.5
1.5 $\text{Mg}(\text{NO}_3)_2$	1				
1.5 $\text{Sr}(\text{NO}_3)_2$	"	29.9912	6.9460	24.1452	7.4

Table 90.3. 1977-1978. 1977-1978

<u>Year</u>	<u>1977</u>	<u>1978</u>
1977.1	1	1977.1 1.1
1977.2	1	1977.2 1.1
1977.3	1	1977.3 1.1
1977.4	1	1977.4 1.1
1977.5	1	1977.5 1.1
1977.6	1	1977.6 1.1
1977.7	1	1977.7 1.1
1977.8	1	1977.8 1.1
1977.9	1	1977.9 1.1
1977.10	1	1977.10 1.1
1977.11	1	1977.11 1.1
1977.12	1	1977.12 1.1

Table 90.4. 1977-1978. 1977-1978

<u>Year</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>
1977.1	1977.1 1.1	1977.1 1.1	1977.1 1.1	1977.1 1.1
1977.2	1977.2 1.1	1977.2 1.1	1977.2 1.1	1977.2 1.1
1977.3	1977.3 1.1	1977.3 1.1	1977.3 1.1	1977.3 1.1
1977.4	1977.4 1.1	1977.4 1.1	1977.4 1.1	1977.4 1.1
1977.5	1977.5 1.1	1977.5 1.1	1977.5 1.1	1977.5 1.1
1977.6	1977.6 1.1	1977.6 1.1	1977.6 1.1	1977.6 1.1
1977.7	1977.7 1.1	1977.7 1.1	1977.7 1.1	1977.7 1.1
1977.8	1977.8 1.1	1977.8 1.1	1977.8 1.1	1977.8 1.1
1977.9	1977.9 1.1	1977.9 1.1	1977.9 1.1	1977.9 1.1
1977.10	1977.10 1.1	1977.10 1.1	1977.10 1.1	1977.10 1.1
1977.11	1977.11 1.1	1977.11 1.1	1977.11 1.1	1977.11 1.1
1977.12	1977.12 1.1	1977.12 1.1	1977.12 1.1	1977.12 1.1

Table LIV. CONDUCTIVITY MEASUREMENTS.

n.	m.	P.	ν_c .	k_w .	k.	η .	Λ_w .	α .
0.3 "	0.15 "	1 "	6.666 "	.025242	.013701 .013164	.003025	.012583 .011555	0.678 .573
0.5 Mg 0.5 Si	0.35 "	1 "	4.0000 "	.035522	.021563 .019995	.005946	.011243 .016681	.537 .497
0.7 Mg 0.7 Si	0.35 "	1 "	2.8571 "	.04660	.028613 .025748	.009701	.024231 .020429	.539 .434
0.9 Mg 0.9 Si	0.45 "	1 "	2.2222 "	.051704	.035123 .030645	.014064	.023881 .022329	.500 .378
1.1 Mg 1.1 Si	0.55 "	1 "	1.8182 "	.057068	.041506 .035106	.019526	.032937 .021096	.467 .326
1.3 Mg 1.3 Si	0.65 "	1 "	1.538 "	.060997	.046942 .038936	.024881	.036202 .024786	.447 .324
1.5 Mg 1.5 Si	0.75 "	1 "	1.3333 "	.065738	.052260 .042076	.030548	.039971 .024476	.408 .243

11 12 13 14 15 16 17

18 19 20 21 22 23 24

25 26 27 28 29 30 31

32 33 34 35 36 37 38

39 40 41 42 43 44 45

46 47 48 49 50 51 52

53 54 55 56 57 58 59

60 61 62 63 64 65 66

67 68 69 70 71 72 73

74 75 76 77 78 79 80

Table LVI.

HYDRATES.

n.	P.	m _c .	α.	L _c .		Δ.	Sp. Gr. Cor.	n _c .	
0.5 Mg ₂ NO ₃	1	0.15	0.638	4.231	5.3	1.594	1.2	4.1	32.5
0.3 Sr ₂ NO ₃	"	"	.573	3.992	2.1			1.0	6.6
0.5 Mg	1	0.25	.587	4.042	6.7	2.351	2.5	6.5	26.0
0.5 Sr	"	"	.497	3.707	3.0			2.4	9.6
0.7 Mg	1	0.35	.539	3.864	9.4	3.371	3.2	8.8	25.1
0.7 Sr	"	"	.434	3.476	4.6			3.4	9.6
0.9 Mg	1	0.45	.500	3.718	11.7	4.513	4.2	11.4	25.2
0.9 Sr	"	"	.378	3.264	5.1			4.4	9.8
1.1 Mg	1	0.55	.467	3.596	13.6	5.646	5.3	13.3	24.2
1.1 Sr	"	"	.326	3.073	5.6			4.9	3.8
1.3 Mg	1	0.65	.433	3.472	15.6	6.892	6.3	15.3	22.4
1.3 Sr	"	"	.284	2.915	6.4			5.5	3.5
1.5 Mg	1	0.75	.408	3.377	17.6	8.169	7.4	17.0	22.7
1.5 Sr	"	"	.243	2.763	7.2			5.7	7.6

The values of n for this mixture are calculated by the method given on pages 61, 62. Evidently, the same method of calculation applies here as in the case of a mixture of calcium chloride and strontium chloride.


IVT eldnt

.M	.I	.	.m	.P	.m
3.2	133.4	833.0	31.0	I	01334 3.0
1.3	322.3	373.	"	"	01312 3.0
7.3	340.4	733.	33.0	I	34 3.0
0.3	707.3	724.	"	"	12 3.0
4.2	433.3	933.	33.0	I	34 7.0
3.4	074.3	434.	"	"	12 7.0
7.11	817.3	003.	34.0	I	34 3.0
1.2	403.3	873.	"	"	12 3.0
3.31	393.3	734.	33.0	I	34 1.1
0.2	370.3	333.	"	"	12 1.1
3.31	374.3	334.	33.0	I	34 3.1
4.3	313.3	433.	"	"	12 3.1
3.71	773.3	804.	37.0	I	34 3.1
3.7	337.3	343.	"	"	12 3.1

The values of M in single solution and in the mixture, for magnesium nitrate and strontium nitrate, have been plotted in curves in Figure VII. The phenomena are, in general, the same as those already pointed out in preceding cases. Curve IV deviates from Curve III with increasing concentration, showing the effect of the large increase in the value of M for magnesium nitrate with increasing concentration, upon the amount of water eliminated as water of hydration by the strontium nitrate. The strontium nitrate is dehydrated, partially, by the magnesium nitrate.

MAGNESIUM NITRATE ALONE.
 " " MIXED WITH STRONTIUM NITRATE.
 I. STRONTIUM NITRATE ALONE.
 " " MIXED WITH MAGNESIUM NITRATE.

DEPT. EXPERIMENTAL ENGINEERING, SIBLEY COLLEGE, CORNELL UNIVERSITY.



GRAM MOLECULES OF SALT PER LITER OF SOLUTION.

FIGURE VII.

In Tables LVII and LVIII, we have the differences in the amount of solvent present and in the values of M , as previously described for the corresponding tables.

Table LVII. MAGNESIUM NITRATE IN MIXTURE OF MAGNESIUM NITRATE AND STRONTIUM NITRATE.

m .	n_{H_2O} .	D_M .
0.15	25	0.4
0.25	56	0.3
0.35	80	0.2
0.45	103	0.3
0.55	118	0.3
0.65	176	0.4
0.75	146	0.6

Table LVIII. STRONTIUM NITRATE IN MIXTURE OF MAGNESIUM NITRATE AND STRONTIUM NITRATE.

m .	n_{H_2O} .	D_M .
0.15	93	1.1
0.25	128	0.6
0.35	172	1.3
0.45	224	0.7
0.55	264	0.7
0.65	304	0.9
0.75	341	1.5

In Figures VII and IX these values are plotted in curves.

2. The first 1000 values of $Y(t)$ are used to estimate the

parameters of the model. The remaining 1000 values are used to

estimate the parameters of the model.

The results of the simulation are shown in Figure 1.

TABLE 1		
Parameters of the model		
α	β	γ
0.1	0.1	0.1
0.2	0.2	0.2
0.3	0.3	0.3
0.4	0.4	0.4
0.5	0.5	0.5
0.6	0.6	0.6
0.7	0.7	0.7
0.8	0.8	0.8
0.9	0.9	0.9
1.0	1.0	1.0


The results of the simulation are shown in Figure 1.

TABLE 2		
Parameters of the model		
α	β	γ
0.1	0.1	0.1
0.2	0.2	0.2
0.3	0.3	0.3
0.4	0.4	0.4
0.5	0.5	0.5
0.6	0.6	0.6
0.7	0.7	0.7
0.8	0.8	0.8
0.9	0.9	0.9
1.0	1.0	1.0

The results of the simulation are shown in Figure 1.

DIFFERENCE BETWEEN WATER PRESENT AS SOLVENT IN THE SINGLE SOLUTION OF MAGNESIUM NITRATE AND IN MIXTURE OF MAGNESIUM NITRATE AND STRONTIUM NITRATE.
 DIFFERENCE BETWEEN VALUES OF α FOR MAGNESIUM NITRATE IN SINGLE SOLUTION AND IN THE MIXTURE OF MAGNESIUM NITRATE AND STRONTIUM NITRATE.

DEPT. EXPERIMENTAL ENGINEERING, SIBLEY COLLEGE, CORNELL UNIVERSITY.




GRAM MOLECULES OF SALT PER LITER OF SOLUTION.
 FIGURE VIII.

DIFFERENCE BETWEEN THE AMOUNT OF WATER PRESENT AS SOLVENT IN THE SINGLE SOLUTION OF STRONTIUM NITRATE AND IN THE MIXTURE OF MAGNESIUM NITRATE AND STRONTIUM NITRATE.

DIFFERENCE BETWEEN THE VALUES OF M FOR STRONTIUM NITRATE IN SINGLE SOLUTION AND IN THE MIXTURE OF MAGNESIUM NITRATE AND STRONTIUM NITRATE.

DEPT. EXPERIMENTAL ENGINEERING, RILEY COLLEGE, CORNELL UNIVERSITY.

I. GRAMS OF WATER.



II. GRAM MOLECULES OF WATER.

GRAM MOLECULES OF SALT PER LITER OF SOLUTION.

FIGURE IX.

We have here, again, a uniform decrease in the amount of water present as solvent, showing a uniform increase in the hydration with increase in concentration. The curves showing the differences in the values of Δ correspond to a constant difference in the amount of water eliminated in single solution and in the mixture. This constant difference is, of course, due to the fact that the concentration of both salts in the mixture is increased uniformly. The curves show, also, that this difference is slightly affected by the difference in the amount of water present as solvent, since they show a tendency to increase as the diminution in water present as solvent becomes greater.

We next took up a Pair of Binary Electrolytes, LITHIUM BROMIDE and SODIUM BROMIDE, Neither of which has a very large Hydrating Power, the Hydrating Power of the Sodium Bromide being somewhat less than that of the Lithium Bromide.

SODIUM BROMIDE.

Table LIX. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ/m .
0.35	0.895	2.58
0.55	1.349	2.57
0.45	1.001	2.56
0.55	1.296	2.54
0.65	1.398	2.54
0.75	1.691	2.59
0.85	2.054	2.59
0.95	2.435	2.62

Table LX. CONDUCTIVITY MEASUREMENTS.

$$\Lambda_{\infty}, 20^{\circ}, = 57.7.$$

m.	v.	k.	Λ_v .	α .
0.25	1.0000	.014508	57.2	0.845
0.35	2.8571	.019628	56.1	.828
0.45	2.3222	.024843	55.2	.815
0.55	1.8182	.029744	54.1	.799
0.65	1.538	.034265	52.7	.779
0.75	1.3333	.039334	52.4	.774
0.85	1.1765	.044909	51.8	.765
0.95	1.0526	.048653	51.4	.760
1.05	0.9523	.053228	50.7	.747

Table LXI. WEIGHT-NORMAL CORRECTIONS.

m.	W _{sol.}	W _{salt.}	W _{H₂O.}	Correction Per Cent.
0.25	25.4629	0.6438	24.8191	0.4
0.35	25.7000	0.9014	24.7987	0.8
0.45	25.8222	1.1589	24.7254	1.1
0.55	26.0735	1.4164	24.6571	1.4
0.65	26.2500	1.6736	24.5873	1.7
0.75	26.4545	1.9314	24.5221	1.9
0.85	26.6155	2.1890	24.4265	2.3
0.95	26.8200	2.4465	24.3735	2.5
1.05	27.0072	2.7040	24.3032	2.8

COMPARISON TABLE

1111

$$f(x) = x^2 - 1$$

x	$f(x)$	$f'(x)$	$f''(x)$	$f'''(x)$
1.0	0.00	2.00	0.00	0.00
1.1	0.21	2.20	0.00	0.00
1.2	0.44	2.40	0.00	0.00
1.3	0.69	2.60	0.00	0.00
1.4	0.96	2.80	0.00	0.00
1.5	1.25	3.00	0.00	0.00
1.6	1.56	3.20	0.00	0.00
1.7	1.89	3.40	0.00	0.00
1.8	2.24	3.60	0.00	0.00
1.9	2.61	3.80	0.00	0.00
2.0	3.00	4.00	0.00	0.00

COMPARISON TABLE

1111

x	$f(x)$	$f'(x)$	$f''(x)$	$f'''(x)$
1.0	1.00	2.00	0.00	0.00
1.1	1.21	2.20	0.00	0.00
1.2	1.44	2.40	0.00	0.00
1.3	1.69	2.60	0.00	0.00
1.4	1.96	2.80	0.00	0.00
1.5	2.25	3.00	0.00	0.00
1.6	2.56	3.20	0.00	0.00
1.7	2.89	3.40	0.00	0.00
1.8	3.24	3.60	0.00	0.00
1.9	3.61	3.80	0.00	0.00
2.0	4.00	4.00	0.00	0.00

Table LVII.		HYDRATES.				
$\frac{m.}{\alpha}$	α	L.	Δ/m	L'.	M.	H.
0.25	0.845	3.47	3.53	3.55	1.0	7.0
0.35	.822	3.40	3.57	3.54	2.2	6.8
0.45	.815	3.38	3.56	3.52	2.2	4.9
0.55	.799	3.35	3.54	3.49	2.2	4.0
0.65	.779	3.31	3.54	3.48	2.7	4.2
0.75	.774	3.30	3.59	3.53	3.5	4.6
0.85	.765	3.28	3.59	3.51	3.6	4.2
0.95	.760	3.27	3.62	3.53	4.1	4.3
1.05	.747	3.25	3.66	3.56	4.8	4.6

LITHIUM BROMIDE.

Table LVIII.	FREEZING-POINT MEASUREMENTS.	
$\frac{m.}{\Delta}$	Δ	Δ/m
0.25	0.917	3.67
0.35	1.292	3.69
0.45	1.679	3.75
0.55	2.050	3.80
0.65	2.314	3.87
0.75	2.957	3.94
0.85	3.418	4.02
0.95	3.905	4.11
1.05	4.281	4.17

Table LIV. CONDUCTIVITY MEASUREMENTS.

n.	v.	k.	Λ_v .	α .
0.35	4.0000	.012099	50.8	.848
0.35	2.8571	.017888	49.4	.825
0.45	2.3322	.021094	48.2	.805
0.55	1.8182	.025855	47.0	.785
0.65	1.538	.029868	45.9	.757
0.75	1.3333	.033996	45.7	.745
0.85	1.1765	.037904	44.6	.734
0.95	1.0526	.04176	44.0	.723
1.05	0.9523	.045505	43.3	

Table LV. WEIGHT-NORMAL CORRECTIONS.

m.	$W_{Sol.}$	$W_{Salt.}$	$W_{H_2O.}$	Correction Per Cent.
0.35	25.7913	0.5457	24.8470	0.6
0.35	25.5641	0.7612	24.8029	0.8
0.45	25.7225	0.9786	24.7459	1.0
0.55	25.8690	1.1961	24.6729	1.3
0.65	26.0558	1.4186	24.6222	1.5
0.75	26.1565	1.6311	24.5274	1.9
0.85	26.2303	1.8495	24.4818	2.1
0.95	26.4558	2.0060	24.7898	2.4
1.05	26.6275	2.2875	24.5400	2.6

Table LVII.

HYDRATES.

m.	α .	L.	Δ/n .	L'.	m.	n.
0.25	0.848	3.44	3.67	3.65	8.2	12.6
0.35	.825	3.79	3.69	3.68	8.7	12.6
0.45	.805	3.76	3.75	3.71	8.8	11.6
0.55	.785	3.82	3.80	3.75	8.1	11.6
0.65	.767	3.29	3.87	3.81	7.5	11.5
0.75	.757	3.27	3.94	3.87	8.6	11.5
0.85	.745	3.25	4.02	3.94	9.7	11.4
0.95	.734	3.23	4.11	4.01	10.8	11.4
1.05	.723	3.21	4.17	4.06	11.6	11.0

MIXTURE OF LITHIUM BROMIDE AND SODIUM BROMIDE.

Table LVIII.

FREEZING-POINT MEASUREMENTS.

m.	P.	Δ .
0.5 LiBr	1	1.833
0.5 NaBr	"	
0.7 LiBr	1	2.596
0.7 NaBr	"	
0.9 LiBr	1	3.415
0.9 NaBr	"	
1.1 LiBr	1	4.325
1.1 NaBr	"	
1.3 LiBr	1	5.284
1.3 NaBr	"	
1.5 LiBr	1	6.297
1.5 NaBr	"	
1.7 LiBr	1	7.298
1.7 NaBr	"	
1.9 LiBr	1	8.399
1.9 NaBr	"	
2.1 LiBr	1	9.539
2.1 NaBr	"	

Table LVIII.

CONDUCTIVITY MEASUREMENTS.

m.	P.	m _c .	v _c .	k _w	k.	D.	k _c .	Λ .	α .
0.5 LiBr	1	0.25	4.0000	.035489	.012699	.001513	.011986	47.9	0.850
0.5 NaBr	"	"	"	"	.014798	"	.017593	51.0	.793
0.7 Li	1	0.55	2.9571	.034327	.017238	.003589	.016073	45.9	.767
0.7 Na	"	"	"	"	.019628	"	.019354	52.2	.770
0.9 Li	1	0.45	2.9222	.042727	.011694	.003810	.014905	44.2	.738
0.9 Na	"	"	"	"	.024843	"	.023322	50.7	.749
1.1 Li	1	0.55	1.8182	.051105	.035855	.004194	.023745	43.2	.721
1.1 Na	"	"	"	"	.039744	"	.027360	49.8	.735
1.3 Li	1	0.65	1.538	.058945	.039863	.005187	.027248	42.2	.704
1.3 Na	"	"	"	"	.034269	"	.031517	48.5	.716
1.5 Li	1	0.75	1.3553	.066486	.027904	.006834	.030769	41.1	.685
1.5 Na	"	"	"	"	.041009	"	.035018	47.6	.703
1.7 Li	1	0.85	1.1765	.078132	.04176	.008781	.037841	39.8	.665
1.7 Na	"	"	"	"	.048853	"	.039991	46.2	.633
1.9 Li	1	0.95	1.0520	.080414	.045505	.010199	.03662	38.8	.648
1.9 Na	"	"	"	"	.053228	"	.044531	45.8	.677
2.1 Li	1	1.05	0.9523	.086247	.073996*	.012086	.039822	37.9	.642
2.1 Na	"	"	"	"	.039734	"	.040815	44.6	.659
	1								

Study 1

Since lithium bromide and sodium bromide are each binary, and dissociated to almost exactly the same extent at equal concentrations of the solutions, the loss in conductivity is apportioned to an equal driving back in the dissociation of the lithium bromide and sodium bromide, except in the mixtures $\left. \begin{smallmatrix} 1.7 \text{ LiBr} \\ 1.7 \text{ NaBr} \end{smallmatrix} \right\}$ and $\left. \begin{smallmatrix} 1.9 \text{ LiBr} \\ 1.9 \text{ NaBr} \end{smallmatrix} \right\}$, where the slight difference in the dissociation of the separate solutions is taken into account.

Table LXIX. WEIGHT-NORMAL CORRECTIONS.

m.	P.	W _{Sol.}	W _{Salt.}	W _{H₂O.}	Correction Per Cent.
0.5 LiBr	1	25.9003	1.18748	24.7128	1.1
0.5 NaBr	"				
0.7 LiBr	1	26.3462	1.6635	24.5877	1.7
0.7 NaBr	"				
0.9 LiBr	1	26.6053	2.1775	24.4678	2.1
0.9 NaBr	"				
1.1 LiBr	1	26.9317	2.6125	24.3192	2.7
1.1 NaBr	"				
1.3 LiBr	1	27.2984	3.0862	24.2121	3.2
1.3 NaBr	"				
1.5 LiBr	1	27.5978	3.5625	24.0852	3.9
1.5 NaBr	"				
1.7 LiBr	1	27.9400	4.0375	23.9025	4.4
1.7 NaBr	"				
1.9 LiBr	1	28.2810	4.5125	23.7765	4.9
1.9 NaBr	"				
2.1 LiBr	1	28.6065	4.9875	23.6190	5.5
2.1 NaBr	"				

The values of W and H for this mixture are calculated

Table LXX.

HYDRATES.

m.	P.	m.c.	α .	L _c .	H.	Δ .	Sp. Gr. Cor.	Me.	He.
0.5 LiBr	1	0.25	.800	3.740	3.2	1.823	1.1	2.8	11.2
0.5 NaBr	"	"	.798	3.344	1.9			1.3	4.8
0.7 LiI	1	0.55	.767	3.286	3.7	2.596	1.7	3.5	10.0
0.7 NaI	"	"	.770	3.293	2.2			2.0	5.7
0.9 LiI	1	0.45	.738	3.233	5.2	3.415	3.1	5.2	11.5
0.9 NaI	"	"	.749	3.253	2.2			2.0	7.5
1.1 LiI	1	0.55	.721	3.301	6.4	4.525	2.7	6.7	13.2
1.1 NaI	"	"	.735	3.227	2.2			2.7	4.9
1.3 LiI	1	0.65	.704	3.107	7.5	5.284	3.2	8.0	12.3
1.3 NaI	"	"	.716	3.192	2.7			2.0	4.5
1.5 LiI	1	0.75	.685	3.135	3.6	6.293	3.9	3.9	11.3
1.5 NaI	"	"	.703	3.168	3.5			3.6	4.8
1.7 LiI	1	0.85	.66	3.096	9.7	7.328	4.4	10.3	13.0
1.7 NaI	"	"	.683	3.13	3.6			3.8	4.4
1.9 LiI	1	0.95	.648	3.065	10.8	8.399	4.9	11.0	11.6
1.9 NaI	"	"	.677	3.119	4.1			4.3	4.4
2.1 LiI	1	1.05	.637	3.032	11.6	9.539	5.5	11.6	11.0
2.1 NaI	"	"	.659	3.085	4.8			4.8	4.6

upon the same basis as that previously employed in the mixture of magnesium nitrate and strontium nitrate. The values thus obtained for the mixture of lithium bromide and sodium bromide are given in the following table.

1. 1. 1. 1. 1. 1. 1. 1. 1. 1.

2. 2. 2. 2. 2. 2. 2. 2. 2. 2.

3. 3. 3. 3. 3. 3. 3. 3. 3. 3.

4. 4. 4. 4. 4. 4. 4. 4. 4. 4.

5. 5. 5. 5. 5. 5. 5. 5. 5. 5.

6. 6. 6. 6. 6. 6. 6. 6. 6. 6.

7. 7. 7. 7. 7. 7. 7. 7. 7. 7.

8. 8. 8. 8. 8. 8. 8. 8. 8. 8.


9. 9. 9. 9. 9. 9. 9. 9. 9. 9.

10. 10. 10. 10. 10. 10. 10. 10. 10. 10.

- I. LITHIUM BROMIDE ALONE.
- II. " " MIXED WITH SODIUM BROMIDE.
- III. SODIUM BROMIDE ALONE.
- IV. " " MIXED WITH LITHIUM BROMIDE.

DEPT. EXPERIMENTAL ENGINEERING, SIBLEY COLLEGE, CORNELL UNIVERSITY.

NUMBER OF GRAM MOLECULES OF WATER COMBINED.



R. C. CARPENTER, STAMPA R. T.

GRAM MOLECULES OF SALT PER LITRE OF SOLUTION.

FIGURE V.

Curve II, representing the value of M in the mixture of lithium bromide and sodium bromide, crosses curve I, the value of M for lithium bromide in single solution. The same phenomenon presents itself in the case of sodium bromide, at a somewhat greater concentration. In an earlier part of this paper it was suggested that the hydrating power of the ions alone, and of these ions when combined to form molecules, would probably be different. If, now, the hydrating power of the lithium bromide and the sodium bromide molecules is greater than that of the respective ions, we might expect that the effect of the smaller amount of water present as solvent in the mixture would be overcome by the greater hydrating power of the undissociated molecules. Evidently, for this increased hydrating power of the molecules over the ions to show itself in the resulting curve, the other salt in the mixture must have little hydrating power, and a comparatively small molecular weight, for if the molecular weight is large, so much solvent will be displaced when the solutions are made up volume normal, that the decreased amount of solvent will obscure any change in hydration due to other causes. Also, the difference in the hydrating power of the ions and molecules should be considerable, in order to overcome the effect

of decrease in amount of solvent free, on a 1% increase in concentration.

The difference between the amount of water acting as solvent towards each salt, in the mixture and in single solution, and the differences between the values of H in the same cases are given in the following tables. A negative value of D for H indicates that the number of grams of water eliminated by a salt in a mixture is greater, instead of less than in single solution.

Table LXII. LITHIUM LEONIDE IN MIXTURES OF LITHIUM BROMIDE AND SODIUM BROMIDE.

M.	D_{H_2O} .	D_H .
0.25	27	0.4
0.35	45	0.2
0.45	47	0
0.55	55	-0.3
0.65	69	-0.5
0.75	85	-0.3
0.85	91	-0.5
0.95	101	-0.3
1.05	115	0

THE UNIVERSITY OF CHICAGO LIBRARY
540 EAST 57TH STREET
CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO LIBRARY
540 EAST 57TH STREET
CHICAGO, ILL. 60637
THE UNIVERSITY OF CHICAGO LIBRARY
540 EAST 57TH STREET
CHICAGO, ILL. 60637
THE UNIVERSITY OF CHICAGO LIBRARY
540 EAST 57TH STREET
CHICAGO, ILL. 60637
THE UNIVERSITY OF CHICAGO LIBRARY
540 EAST 57TH STREET
CHICAGO, ILL. 60637

THE UNIVERSITY OF CHICAGO LIBRARY

540 EAST 57TH STREET

DATE	AMOUNT	REMARKS
1967	100	100.00
1968	100	100.00
1969	100	100.00
1970	100	100.00
1971	100	100.00
1972	100	100.00
1973	100	100.00
1974	100	100.00
1975	100	100.00
1976	100	100.00
1977	100	100.00
1978	100	100.00
1979	100	100.00
1980	100	100.00
1981	100	100.00
1982	100	100.00
1983	100	100.00
1984	100	100.00
1985	100	100.00
1986	100	100.00
1987	100	100.00
1988	100	100.00
1989	100	100.00
1990	100	100.00
1991	100	100.00
1992	100	100.00
1993	100	100.00
1994	100	100.00
1995	100	100.00
1996	100	100.00
1997	100	100.00
1998	100	100.00
1999	100	100.00
2000	100	100.00
2001	100	100.00
2002	100	100.00
2003	100	100.00
2004	100	100.00
2005	100	100.00
2006	100	100.00
2007	100	100.00
2008	100	100.00
2009	100	100.00
2010	100	100.00
2011	100	100.00
2012	100	100.00
2013	100	100.00
2014	100	100.00
2015	100	100.00
2016	100	100.00
2017	100	100.00
2018	100	100.00
2019	100	100.00
2020	100	100.00
2021	100	100.00
2022	100	100.00
2023	100	100.00
2024	100	100.00
2025	100	100.00
2026	100	100.00
2027	100	100.00
2028	100	100.00
2029	100	100.00
2030	100	100.00
2031	100	100.00
2032	100	100.00
2033	100	100.00
2034	100	100.00
2035	100	100.00
2036	100	100.00
2037	100	100.00
2038	100	100.00
2039	100	100.00
2040	100	100.00
2041	100	100.00
2042	100	100.00
2043	100	100.00
2044	100	100.00
2045	100	100.00
2046	100	100.00
2047	100	100.00
2048	100	100.00
2049	100	100.00
2050	100	100.00
2051	100	100.00
2052	100	100.00
2053	100	100.00
2054	100	100.00
2055	100	100.00
2056	100	100.00
2057	100	100.00
2058	100	100.00
2059	100	100.00
2060	100	100.00
2061	100	100.00
2062	100	100.00
2063	100	100.00
2064	100	100.00
2065	100	100.00
2066	100	100.00
2067	100	100.00
2068	100	100.00
2069	100	100.00
2070	100	100.00
2071	100	100.00
2072	100	100.00
2073	100	100.00
2074	100	100.00
2075	100	100.00
2076	100	100.00
2077	100	100.00
2078	100	100.00
2079	100	100.00
2080	100	100.00
2081	100	100.00
2082	100	100.00
2083	100	100.00
2084	100	100.00
2085	100	100.00
2086	100	100.00
2087	100	100.00
2088	100	100.00
2089	100	100.00
2090	100	100.00
2091	100	100.00
2092	100	100.00
2093	100	100.00
2094	100	100.00
2095	100	100.00
2096	100	100.00
2097	100	100.00
2098	100	100.00
2099	100	100.00
2100	100	100.00

Table LXIII. SODIUM BROMIDE IN MIXTURE OF LITHIUM BROMIDE AND SODIUM IODIDE.

$n.$	$n_{H_2O}.$	$n_m.$
0.25	54	0.7
0.35	72	0.2
0.45	104	0.2
0.55	144	-0.1
0.65	159	-0.2
0.75	180	-0.1
0.85	205	-0.2
0.95	222	-0.1
1.05	276	0


These values are plotted in curves in Figures XI and XII. The fact that the curves show positive values of n_m has already been explained as probably due to the increased hydrating power of the molecule over its ions, so that when sufficient driving back in dissociation occurs we would get an increased amount of water eliminated as water of hydration. That this difference is nearly constant is to be expected, since corresponding additions of sodium bromide and lithium bromide are always made, in changes of concentration.

TABLE 1. Results of the analysis of variance for the effect of the concentration of the solution on the rate of the reaction.

Concentration of the solution, g/l	Rate of the reaction, g/h	F
0.1	0.1	0.1
0.2	0.2	0.2
0.3	0.3	0.3
0.4	0.4	0.4
0.5	0.5	0.5
0.6	0.6	0.6
0.7	0.7	0.7
0.8	0.8	0.8
0.9	0.9	0.9
1.0	1.0	1.0


The results of the analysis of variance for the effect of the concentration of the solution on the rate of the reaction are shown in Table 1. It can be seen from the table that the rate of the reaction increases with an increase in the concentration of the solution. The rate of the reaction is highest at a concentration of 1.0 g/l. The results of the analysis of variance for the effect of the concentration of the solution on the rate of the reaction are shown in Table 1. It can be seen from the table that the rate of the reaction increases with an increase in the concentration of the solution. The rate of the reaction is highest at a concentration of 1.0 g/l.

TABLE NO. 1. THE AMOUNT OF WATER PRESENT AS SOLVENT IN THE SINGLE SOLUTION OF LITHIUM BROMIDE AND IN THE MIXTURE OF LITHIUM BROMIDE AND SODIUM BROMIDE. DIFFERENCE BETWEEN THE VALUES OF λ FOR LITHIUM BROMIDE IN SINGLE SOLUTION AND IN THE MIXTURE OF LITHIUM BROMIDE AND SODIUM BROMIDE.



DIFFERENCE BETWEEN THE VALUES OF λ FOR SODIUM BROMIDE IN SINGLE SOLUTION AND IN THE MIXTURE OF SODIUM BROMIDE AND LITHIUM BROMIDE.

DIFFERENCE BETWEEN THE VALUES OF λ FOR SODIUM BROMIDE IN SINGLE SOLUTION AND IN THE MIXTURE OF SODIUM BROMIDE AND LITHIUM BROMIDE.



GRAM MOLECULES OF SALT PER LITER OF SOLUTION.
FIGURE XII.

We next consider a Mixture of Calcium Nitrate and Magnesium Nitrate, Both of which Have Hydrating Power.

CALCIUM NITRATE.

Table LXXIII. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ/m .
0.35	1.9189	4.756
0.45	2.094	4.65
0.65	3.041	4.68
0.85	4.057	4.77
1.05	5.123	4.88
1.25	6.345	5.08
1.45	7.547	5.21
1.6555	8.859	5.35

Table LXXIV. CONDUCTIVITY MEASUREMENTS.

m.	v.	k.	Λ_v .	α .
0.25	4.0000	.020975	83.9	0.624
0.45	2.2222	.033259	73.9	.550
0.65	1.538	.042932	66.0	.491
0.85	1.1765	.050272	59.2	.440
1.05	0.9525	.056301	53.6	.399
1.25	0.8000	.060363	48.3	.359
1.45	0.6897	.063236	43.6	.324
1.6555	0.6041	.064674	39.1	.291

and the results are listed in Table 1. The results are given in

Table 1. The results are given in Table 1. The results are given in

Table 1. The results are given in Table 1. The results are given in

Table 1. The results are given in Table 1. The results are given in

Δ	Δ	Δ
10.1	10.1	10.1
10.2	10.2	10.2
10.3	10.3	10.3
10.4	10.4	10.4
10.5	10.5	10.5
10.6	10.6	10.6
10.7	10.7	10.7
10.8	10.8	10.8
10.9	10.9	10.9
11.0	11.0	11.0

Table 1. The results are given in Table 1. The results are given in

Δ	Δ	Δ	Δ	Δ
10.1	10.1	10.1	10.1	10.1
10.2	10.2	10.2	10.2	10.2
10.3	10.3	10.3	10.3	10.3
10.4	10.4	10.4	10.4	10.4
10.5	10.5	10.5	10.5	10.5
10.6	10.6	10.6	10.6	10.6
10.7	10.7	10.7	10.7	10.7
10.8	10.8	10.8	10.8	10.8
10.9	10.9	10.9	10.9	10.9
11.0	11.0	11.0	11.0	11.0

Table LXXV. WEIGHT-NORMAL CORRECTIONS.

m.	W _{Sol.}	W _{Salt.}	W _{H₂O.}	Correction Per Cent.
0.25	25.77665	1.026	24.7405	1.0
0.45	26.3748	1.8467	24.5281	1.9
0.65	26.969"	2.6674	24.3019	2.8
0.85	27.5178	3.4883	24.0286	3.9
2.05	28.1060	4.3089	23.7971	4.8
1.25	28.6822	5.1297	23.5525	5.8
1.45	29.2340	5.9504	23.2836	6.9
1.6555	29.8480	6.7938	23.0542	7.8

Table LXXVI. HYDRATES.

m.	α .	L.	Δ /m.	L'.	M.	H.
0.25	0.624	4.18	4.76	4.71	6.3	25.2
0.45	.550	3.90	4.65	4.56	8.0	17.9
0.65	.491	3.69	4.68	4.55	10.5	16.2
0.85	.440	3.50	4.77	4.58	13.1	15.4
1.05	.391	3.34	4.88	4.65	15.7	15.0
1.25	.359	3.20	5.08	4.79	18.4	14.7
1.45	.324	3.07	5.21	4.85	20.4	14.1
1.6555	.291	2.94	5.35	4.93	22.4	13.5

MAGNESIUM NITRATE.

TABLE 1 Summary of data for the 1941-1942 season

<u>Station</u>	<u>Depth</u>	<u>Time</u>	<u>Wind</u>	<u>Temp</u>
1	1000	1000	1000	1000
2	1000	1000	1000	1000
3	1000	1000	1000	1000
4	1000	1000	1000	1000
5	1000	1000	1000	1000
6	1000	1000	1000	1000
7	1000	1000	1000	1000
8	1000	1000	1000	1000
9	1000	1000	1000	1000
10	1000	1000	1000	1000

TABLE 2

Summary of data for the 1941-1942 season

<u>Station</u>	<u>Depth</u>	<u>Time</u>	<u>Wind</u>	<u>Temp</u>
1	1000	1000	1000	1000
2	1000	1000	1000	1000
3	1000	1000	1000	1000
4	1000	1000	1000	1000
5	1000	1000	1000	1000
6	1000	1000	1000	1000
7	1000	1000	1000	1000
8	1000	1000	1000	1000
9	1000	1000	1000	1000
10	1000	1000	1000	1000

Continued on next page

Table LXXVII. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ/m .
0.25	1.252	5.01
0.45	2.795	5.52
0.65	3.669	5.64
0.85	5.113	6.02
1.05	6.762	6.44
1.25	8.562	6.85
1.45	10.589	7.30

Table LXXVIII. CONDUCTIVITY MEASUREMENTS.

$$\Lambda_{\infty}, 0^{\circ}, = 128.5.$$

m.	v.	k.	Λ_v .	α 0.671.
0.25	4.0000	.021565	86.3	
0.45	2.2222	.075123	78.1	.607
0.65	1.538	.046942	72.2	.562
0.85	1.1765	.056821	66.9	.520
1.05	0.9525	.065334	62.2	.484
1.25	0.8000	.071668	57.7	.446
1.45	0.68966	.076279	52.6	.409

Table LXXIX. WEIGHT-NORMAL CORRECTIONS.

m.	Wsol.	Wsalt.	WH ₂ O.	Correction Per Cent.
0.25	25.648	0.9277	24.7008	1.0
0.45	26.2075	1.6699	24.5376	1.8
0.65	26.7605	2.4122	24.7483	2.6
0.85	27.1958	3.1543	24.0415	3.8
1.05	27.7080	3.8965	23.8115	4.8
1.25	28.2455	4.6287	23.6048	5.6
1.45	28.7626	5.3809	23.7817	6.5

Table LXXX. HYDRATES.

m.	α .	L.	Δ/m .	L'.	M.	H.
0.25	0.671	4.86	5.01	4.96	6.7	26.2
0.45	.607	4.12	5.32	5.22	11.7	26.0
0.65	.562	3.95	5.64	5.49	15.6	24.0
0.85	.520	3.80	6.02	5.79	19.1	22.5
1.05	.484	3.66	6.44	6.13	22.4	21.3
1.25	.446	3.52	6.85	6.47	25.3	20.2
1.45	.409	3.38	7.30	6.83	28.1	19.4

Data for the Mixture of CALCIUM NITRATE and MAGNESIUM NITRATE follows.

TABLE 1. SUMMARY OF DATA FOR THE STUDY

Year	Age	Sex	Weight (kg)	Height (cm)	Length (cm)	Area (cm ²)
1990	10	M	25.0	120.0	18.0	150.0
1991	11	F	28.0	125.0	19.0	160.0
1992	12	M	30.0	130.0	20.0	170.0
1993	13	F	32.0	135.0	21.0	180.0
1994	14	M	35.0	140.0	22.0	190.0
1995	15	F	38.0	145.0	23.0	200.0
1996	16	M	40.0	150.0	24.0	210.0
1997	17	F	42.0	155.0	25.0	220.0

TABLE 2. SUMMARY OF DATA FOR THE STUDY

Year	Age	Sex	Weight (kg)	Height (cm)	Length (cm)	Area (cm ²)
1998	18	M	45.0	160.0	26.0	230.0
1999	19	F	48.0	165.0	27.0	240.0
2000	20	M	50.0	170.0	28.0	250.0
2001	21	F	52.0	175.0	29.0	260.0
2002	22	M	55.0	180.0	30.0	270.0
2003	23	F	58.0	185.0	31.0	280.0
2004	24	M	60.0	190.0	32.0	290.0
2005	25	F	62.0	195.0	33.0	300.0
2006	26	M	65.0	200.0	34.0	310.0
2007	27	F	68.0	205.0	35.0	320.0

NOTE: Data were collected from the study and are presented in the table.

Source: Author's calculations.

Table LXXII. FREEZING-POINT MEASUREMENTS.

m.	P.	Δ .
0.5 $\text{Ca}(\text{NO}_3)_2$	1	29.492
0.5 $\text{Mg}(\text{NO}_3)_2$	"	
0.9 $\text{Ca}(\text{NO}_3)_2$	1	4.856
0.9 $\text{Mg}(\text{NO}_3)_2$	"	
1.3 $\text{Ca}(\text{NO}_3)_2$	1	7.624
1.3 $\text{Mg}(\text{NO}_3)_2$	"	
1.7 $\text{Ca}(\text{NO}_3)_2$	1	10.95
1.7 $\text{Mg}(\text{NO}_3)_2$	"	
2.1 $\text{Ca}(\text{NO}_3)_2$	1	19.90
2.1 $\text{Mg}(\text{NO}_3)_2$	"	
2.5 $\text{Ca}(\text{NO}_3)_2$	1	19.73
2.5 $\text{Mg}(\text{NO}_3)_2$	"	
2.9 $\text{Ca}(\text{NO}_3)_2$	1	24.8
2.9 $\text{Mg}(\text{NO}_3)_2$	"	

Table LXXIII. WEIGHT-NORMAL CORRECTIONS.

m.	P.	$W_{\text{Sol.}}$	$W_{\text{Salt.}}$	$W_{\text{H}_2\text{O.}}$	Correction Per Cent.
0.5 $\text{Ca}(\text{NO}_3)_2$	1	26.4387	1.9557	24.4850	2.1
0.5 $\text{Mg}(\text{NO}_3)_2$	"				
0.9 $\text{Ca}(\text{NO}_3)_2$	1	27.5700	3.5166	24.0134	3.9
0.9 $\text{Mg}(\text{NO}_3)_2$	"				
1.3 $\text{Ca}(\text{NO}_3)_2$	1	28.5974	5.0796	23.5178	5.9
1.3 $\text{Mg}(\text{NO}_3)_2$	"				
1.7 $\text{Ca}(\text{NO}_3)_2$	1	29.6600	6.6425	23.0175	7.9
1.7 $\text{Mg}(\text{NO}_3)_2$	"				
2.1 $\text{Ca}(\text{NO}_3)_2$	1	30.6258	8.2055	22.4193	10.3
2.1 $\text{Mg}(\text{NO}_3)_2$	"				
2.5 $\text{Ca}(\text{NO}_3)_2$	1	31.6538	9.7684	21.8854	12.5
2.5 $\text{Mg}(\text{NO}_3)_2$	"				
2.9 $\text{Ca}(\text{NO}_3)_2$	1	32.6175	11.3315	21.2862	14.9
2.9 $\text{Mg}(\text{NO}_3)_2$	"				

ANNUAL REPORT 2023

	2023	2022
Revenue	1,234,567	1,123,456
Expenses	876,543	765,432
Profit	358,024	358,024
Assets	5,678,901	5,678,901
Liabilities	2,345,678	2,345,678
Equity	3,333,223	3,333,223

FINANCIAL STATEMENTS

Item	2023	2022	2021	2020
Revenue	1,234,567	1,123,456	1,012,345	901,234
Expenses	876,543	765,432	654,321	543,210
Profit	358,024	358,024	358,024	358,024
Assets	5,678,901	5,678,901	5,678,901	5,678,901
Liabilities	2,345,678	2,345,678	2,345,678	2,345,678
Equity	3,333,223	3,333,223	3,333,223	3,333,223

Table LXXVIII.

m.	p.	m_c .	v_c .	k_w .	k_c .	Λ_v .	α .
0.5 CaO ₃	1	0.35	1.0000	.036833	.030975	.019011	.71
0.5 CaO ₃	"	"	"		.091563	.619320	
0.9 CaO	1	0.45	2.3233	.054700	.053959	.025920	.498
0.9 CaO	"	"	"		.075133	.018790	.498
1.3 CaO	1	0.65	1.533	.066970	.042932	.029904	.542
1.3 CaO	"	"	"		.046942	.070066	.482
1.7 CaO	1	0.85	1.1765	.071201	.050272	.030420	.366
1.7 CaO	"	"	"		.056831	.040781	.73
2.1 CaO	1	1.05	0.9523	.071673	.056301	.028332	.301
2.1 CaO	"	"	"		.065774	.043741	
2.5 CaO	1	1.25	0.8000	.069140	.060767	.024807	.3
2.5 CaO	"	"	"		.071668	.014435	.976
2.9 CaO	1	1.45	0.6966	.067337	.063336	.019866	.302
2.9 CaO	"	"	"		.076279	.043461	.973

四 五 六 七 八 九 十

一 二 三 四 五 六 七 八 九 十

一 二 三 四 五 六 七 八 九 十

一 二 三 四 五 六 七 八 九 十

一 二 三 四 五 六 七 八 九 十

一 二 三 四 五 六 七 八 九 十

一 二 三 四 五 六 七 八 九 十

一 二 三 四 五 六 七 八 九 十

一 二 三 四 五 六 七 八 九 十

一 二 三 四 五 六 七 八 九 十

Table LXXIV.

HYDRATES.

m.	P.	n _c .	α.	l _c .	M.	Δ	Sp.Gr. Cor.	n _c .	H _c .
0.5 Ba(NO ₃) ₂	1	0.25	0.565	3.96	6.3	2.498	2.1	4.6	18.5
0.5 Ba(NO ₃) ₂	"	"	.617	4.16	6.7			5.3	20.8
0.9 Ca	1	0.45	.428	3.45	8.0	4.856	3.9	7.2	16.0
0.9 Mg	"	"	.498	3.71	11.7			11.1	24.7
1.3 Ca	1	0.65	.342	3.13	10.5	7.624	5.9	9.4	14.4
1.3 Mg	"	"	.432	3.47	15.6			14.9	22.9
1.7 Ca	1	0.85	.266	2.85	13.1	10.93	7.9	11.7	13.8
1.7 Mg	"	"	.373	3.25	19.1			18.1	21.3
2.1 Ca	1	1.05	.201	2.61	15.7	14.90	10.3	15.6	13.0
2.1 Mg	"	"	.321	3.05	22.4			21.0	20.0
2.5 Ca	1	1.25	.148	2.41	18.4	19.73	12.5	15.7	12.5
2.5 Mg	"	"	.276	2.89	25.3			25.3	18.7
2.9 Ca	1	1.45	.102	2.24	20.4	27.8	14.9	16.5	11.4
2.9 Mg	"	"	.235	2.75	28.1			25.2	17.4

三 龍 龍 龍 龍 龍 龍 龍

三 龍 龍 龍 龍 龍 龍 龍

三 龍 龍 龍 龍 龍 龍 龍

三 龍 龍 龍 龍 龍 龍 龍

三 龍 龍 龍 龍 龍 龍 龍

三 龍 龍 龍 龍 龍 龍 龍

三 龍 龍 龍 龍 龍 龍 龍

三 龍 龍 龍 龍 龍 龍 龍

三 龍 龍 龍 龍 龍 龍 龍

三 龍 龍 龍 龍 龍 龍 龍

The values of M for each salt in the mixture and in separate solution are plotted in curves in Figure XIII. With increased concentration we have increased divergence in curves I and III which represent the hydrating power (values of M) in separate solutions of magnesium nitrate and of calcium nitrate, respectively. Consequently, as we should expect, curves II and IV, representing the values of M of the two salts in the mixture, show increasing divergence with increase in concentration.



the value of λ is the same as the value of λ in the case of

the case of λ is the same as the value of λ in the case of

the case of λ is the same as the value of λ in the case of

the case of λ is the same as the value of λ in the case of

the case of λ is the same as the value of λ in the case of

the case of λ is the same as the value of λ in the case of

the case of λ is the same as the value of λ in the case of

the case of λ is the same as the value of λ in the case of

the case of λ is the same as the value of λ in the case of


MAGNESIUM NITRATE ALONE.

" " MIXED WITH CALCIUM NITRATE.

CALCIUM NITRATE ALONE.

" " MIXED WITH MAGNESIUM NITRATE.

DEPT. EXPERIMENTAL ENGINEERING, AIRLEY COLLEGE, CORNELL UNIVERSITY.



GRAM MOLECULES OF SALT PER LITRE OF SOLUTION.

FIGURE VIII.

Table LXXXVI. CALCIUM NITRATE IN MIXTURE OF CALCIUM NITRATE AND MAGNESIUM NITRATE.

m.	D_{H_2O} .	D_M .
0.25	104.4	1.7
0.45	220.3	0.8
0.65	298.5	1.1
0.85	365.7	1.4
1.05	432.5	2.1
1.25	487	2.7
1.45	534	3.9

Table LXXXVII MAGNESIUM NITRATE IN MIXTURE OF CALCIUM NITRATE AND MAGNESIUM NITRATE.

m.	D_{H_2O} .	D_M .
0.25	93.7	1.5
0.45	150.5	0.6
0.65	202	0.7
0.85	251	1.0
1.05	300	1.4
1.25	351	2.0
1.45	381	2.9

These values are plotted in curves in Figures XIV and XV.

TABLE 1. The results of the first series of experiments.

TABLE 1. The results of the first series of experiments.

θ	θ_0	θ
0.1	0.01	0.01
0.2	0.02	0.02
0.3	0.03	0.03
0.4	0.04	0.04
0.5	0.05	0.05
0.6	0.06	0.06
0.7	0.07	0.07
0.8	0.08	0.08

TABLE 2. The results of the second series of experiments.

TABLE 2. The results of the second series of experiments.


θ	θ_0	θ
0.1	0.01	0.01
0.2	0.02	0.02
0.3	0.03	0.03
0.4	0.04	0.04
0.5	0.05	0.05
0.6	0.06	0.06
0.7	0.07	0.07
0.8	0.08	0.08

TABLE 3. The results of the third series of experiments.

TABLE 3.

DETERMINED BY THE VALUE OF λ WHICH GIVES THE BEST FIT TO THE EXPERIMENTAL
 DATA OF CALCIUM NITRATE AND OF THE MIXTURE OF CALCIUM NITRATE AND MAGNESIUM
 NITRATE.
 DIFFERENCE BETWEEN THE VALUES OF λ FOR CALCIUM NITRATE IN SEVERAL SOLUTIONS AND
 IN THE MIXTURE OF CALCIUM NITRATE AND MAGNESIUM NITRATE.

DEPT. EXPERIMENTAL ENGINEERING, SIKSKY COLLEGE, CORNELL UNIVERSITY.




B. C. CARPENTER, STANFORD U. N. Y.

GRAM MOLECULARS OF SALT PER LITER OF SOLUTION.
 FIGURE XIV.

GRAMS OF SALT PER LITER OF WATER.

DETERMINED FROM THE VALUES OF μ FOR VARIOUS TEMPERATURES IN THE SYSTEM OF
 SOLUTION OF CALCIUM HYDROXIDE AND IN THE SYSTEM OF MAGNESIUM HYDROXIDE AND CALCIUM
 HYDROXIDE. DETERMINED THE VALUES OF μ FOR VARIOUS TEMPERATURES IN THE SYSTEM OF
 SOLUTION OF CALCIUM HYDROXIDE AND MAGNESIUM HYDROXIDE.

I. GRAMS OF WATER.



II. GRAM MOLES OF WATER.

GRAM MOLES OF SOLUTE PER LITER OF SOLUTION.

FIGURE 17.

The curves in Figures XIV and XV show the same general relations as the preceding curves of the same character, excepting those of lithium bromide and sodium bromide. The amount of water eliminated to form hydrates in the mixture, tends to diminish somewhat proportionally to the diminution in the amount of water present which is acting as solvent. This is clearly shown by the direction in which the curves turn with increased concentration of the solutions. It is probable that in the more dilute solutions we have the difference in the hydrating power of the ions and molecules affecting the total amount of water eliminated as water of hydration. Consequently, where the total amount of salt is constant, but a change in the association is produced, if the number of ions is large relative to the number of molecules unassociated, as is the case in the more dilute solutions, the curve for D_m does not at once show the influence of diminishing amount of solvent present. This phenomenon is shown by the curves as change of direction, or increased curvature in the curve which represents the value of D_m .

A Pair of Quaternary Electrolytes was Next Studied. Ferric chloride and aluminium chloride were employed. Very concentrated stock solutions were prepared, and the necess-

The first is the fact that the
 second is the fact that the
 third is the fact that the
 fourth is the fact that the
 fifth is the fact that the
 sixth is the fact that the
 seventh is the fact that the
 eighth is the fact that the
 ninth is the fact that the
 tenth is the fact that the
 eleventh is the fact that the
 twelfth is the fact that the
 thirteenth is the fact that the
 fourteenth is the fact that the
 fifteenth is the fact that the
 sixteenth is the fact that the
 seventeenth is the fact that the
 eighteenth is the fact that the
 nineteenth is the fact that the
 twentieth is the fact that the
 twenty-first is the fact that the
 twenty-second is the fact that the
 twenty-third is the fact that the
 twenty-fourth is the fact that the
 twenty-fifth is the fact that the
 twenty-sixth is the fact that the
 twenty-seventh is the fact that the
 twenty-eighth is the fact that the
 twenty-ninth is the fact that the
 thirtieth is the fact that the
 thirty-first is the fact that the
 thirty-second is the fact that the
 thirty-third is the fact that the
 thirty-fourth is the fact that the
 thirty-fifth is the fact that the
 thirty-sixth is the fact that the
 thirty-seventh is the fact that the
 thirty-eighth is the fact that the
 thirty-ninth is the fact that the
 fortieth is the fact that the
 forty-first is the fact that the
 forty-second is the fact that the
 forty-third is the fact that the
 forty-fourth is the fact that the
 forty-fifth is the fact that the
 forty-sixth is the fact that the
 forty-seventh is the fact that the
 forty-eighth is the fact that the
 forty-ninth is the fact that the
 fiftieth is the fact that the
 fifty-first is the fact that the
 fifty-second is the fact that the
 fifty-third is the fact that the
 fifty-fourth is the fact that the
 fifty-fifth is the fact that the
 fifty-sixth is the fact that the
 fifty-seventh is the fact that the
 fifty-eighth is the fact that the
 fifty-ninth is the fact that the
 sixtieth is the fact that the
 sixty-first is the fact that the
 sixty-second is the fact that the
 sixty-third is the fact that the
 sixty-fourth is the fact that the
 sixty-fifth is the fact that the
 sixty-sixth is the fact that the
 sixty-seventh is the fact that the
 sixty-eighth is the fact that the
 sixty-ninth is the fact that the
 seventieth is the fact that the
 seventy-first is the fact that the
 seventy-second is the fact that the
 seventy-third is the fact that the
 seventy-fourth is the fact that the
 seventy-fifth is the fact that the
 seventy-sixth is the fact that the
 seventy-seventh is the fact that the
 seventy-eighth is the fact that the
 seventy-ninth is the fact that the
 eightieth is the fact that the
 eighty-first is the fact that the
 eighty-second is the fact that the
 eighty-third is the fact that the
 eighty-fourth is the fact that the
 eighty-fifth is the fact that the
 eighty-sixth is the fact that the
 eighty-seventh is the fact that the
 eighty-eighth is the fact that the
 eighty-ninth is the fact that the
 ninetieth is the fact that the
 ninety-first is the fact that the
 ninety-second is the fact that the
 ninety-third is the fact that the
 ninety-fourth is the fact that the
 ninety-fifth is the fact that the
 ninety-sixth is the fact that the
 ninety-seventh is the fact that the
 ninety-eighth is the fact that the
 ninety-ninth is the fact that the
 hundredth is the fact that the

ary measurements were made as quickly as possible, in order to reduce hydrolysis to a minimum, there being a time factor in the hydrolysis of such compounds as ferric chloride.

Notwithstanding all precautions there was, of course, considerable hydrolysis. Especially was this the case in the mixture of the two salts, the presence of the aluminium chloride appearing to act catalytically upon the solution of ferric chloride, causing a gradual precipitation of hydroxide of iron.

ALUMINIUM CHLORIDE.

Table LXXVIII. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ /m.
0.2	1.279	6.40
0.4	2.910	7.28
0.6	5.144	8.57
0.8	7.962	9.95
1.0	11.795	11.80
1.2	16.385	13.65
1.3617	21.75	15.97
1.415	23.5	16.6

Table LXVIII. CONDUCTIVITY MEASUREMENTS.

m.	v.	k.	Λ_v .	α .
0.2	5.000	.023983	119.9	0.660
0.4	2.500	.04053	101.3	.558
0.6	1.6666	.058115	88.5	.487
0.8	1.250	.06118	76.5	.421
1.0	1.000	.066595	66.6	.367
1.2	0.8333	.06842	57.0	.314
1.3617	0.7344	.06760	49.7	.273

Table XC. WEIGHT-NORMAL CORRECTIONS.

m.	$W_{\text{Sol.}}$	$W_{\text{Salt.}}$	$W_{\text{H}_2\text{O.}}$	Correction Per Cent.
0.2	25.6340	0.6673	24.9567	0.2
0.4	26.1538	1.3346	24.8192	0.7
0.6	26.7134	2.0019	24.7115	1.2
0.8	27.2153	2.6692	24.5461	1.8
1.0	27.7351	3.3365	24.3986	2.4
1.2	28.2977	4.0038	24.2839	2.9
1.3617	28.7252	4.5433	24.1819	3.3

COMPARISON STATISTICS - DATA SET

α	$\sqrt{F_0}$	α	β	α
0.05	1.072	0.0500	1.072	0.2
0.10	1.171	0.0500	1.072	0.5
0.20	1.300	0.1000	1.072	1.0
0.50	1.617	0.5000	1.072	5.0
1.00	1.910	0.9000	1.072	10.0
1.00	2.110	1.0000	1.072	15.0
1.00	2.112	1.0000	1.072	20.0

COMPARISON STATISTICS - 100 DATA

COMPARISON STATISTICS	$\sqrt{F_0}$	α	β	α
0.05	1.072	0.0500	1.072	0.2
0.10	1.171	0.0500	1.072	0.5
0.20	1.300	0.1000	1.072	1.0
0.50	1.617	0.5000	1.072	5.0
1.00	1.910	0.9000	1.072	10.0
1.00	2.110	0.9000	1.072	15.0
1.00	2.112	0.9000	1.072	20.0

Table XCI. HYDRATES.

m.	α .	L.	Δ/m .	L'.	M.	H.
0.2	0.060	5.54	6.40	6.39	7.4	37.0
0.4	.558	4.97	7.28	7.23	17.4	43.5
0.6	.487	4.58	8.57	8.47	25.5	42.5
0.8	.421	4.21	9.95	9.77	31.6	39.5
1.0	.367	3.91	11.80	11.52	36.7	36.7
1.2	.314	3.61	13.65	13.25	40.4	33.7
1.3617	.273	3.39	15.97	15.44	43.4	31.9

FERRIC CHLORIDE.

Table XCII. FREEZING-POINT MEASUREMENTS.

m.	Δ .	Δ/m .
0.2	1.255	6.28
0.4	2.715	6.79
0.6	4.530	7.55
0.8	6.667	8.33
1.0	9.152	9.15
1.2	12.11	10.09
1.3617	14.592	10.72

				CONCENTRATION	TOTAL AREA	
λ	μ	λ/μ	λ/Δ	Δ	λ/δ	μ
1.00	2.00	05.0	100.0	10.0	900.0	9.0
1.00	2.50	04.0	125.0	12.5	800.0	8.0
1.00	3.00	33.3	150.0	15.0	700.0	7.0
1.00	3.50	28.6	175.0	17.5	600.0	6.0
1.00	4.00	25.0	200.0	20.0	500.0	5.0
1.00	4.50	22.2	225.0	22.5	400.0	4.0
1.00	5.00	20.0	250.0	25.0	360.0	3.6
1.00	5.50	18.2	275.0	27.5	330.0	3.3
1.00	6.00	16.7	300.0	30.0	300.0	3.0

CONCENTRATION RATIO

CONCENTRATION RATIO	CONCENTRATION RATIO	TOTAL AREA
λ/Δ	Δ	μ
100.0	10.0	9.0
125.0	12.5	8.0
150.0	15.0	7.0
175.0	17.5	6.0
200.0	20.0	5.0
225.0	22.5	4.0
250.0	25.0	3.6
275.0	27.5	3.3
300.0	30.0	3.0

Table XCIII. CONDUCTIVITY MEASUREMENTS.

$$\Lambda_{\infty, 0^{\circ}} = 220. \quad (\text{Calculated}).$$

m.	v.	k.	Λ_v .	α .
0.2	5.000	.02380	119.0	0.541
0.4	2.500	.03786	94.7	.430
0.6	1.6666	.04645	77.4	.352
0.8	1.250	.05147	64.3	.292
1.0	1.000	.053147	53.1	.241
1.2	0.8333	.053145	44.3	.201
1.3617	0.7344	.051777	38.0	.173

Table XCIV. WEIGHT-NORMAL CORRECTIONS.

m.	W _{Sol.}	W _{Salt.}	W _{H₂O.}	Correction Per Cent.
0.2	25.6792	0.8113	24.8679	0.5
0.4	26.3380	1.6225	24.7155	1.1
0.6	26.9956	2.4337	24.5619	1.8
0.8	27.6250	3.245	24.378	2.5
1.0	28.2400	4.056	24.184	3.3
1.2	28.8367	4.8675	24.0192	3.9
1.3617	29.3690	5.5234	23.8456	4.6

TABLE 10. *Estimated values of β_{max} and $\beta_{\text{max}}/\beta_{\text{min}}$ for various values of β_{min} and β_{max}*

β_{min}	β_{max}	β_{max}	β_{max}	β_{max}
0.01	0.02	0.0100	0.0100	1.0
0.02	0.04	0.0200	0.0200	1.0
0.03	0.06	0.0300	0.0300	1.0
0.04	0.08	0.0400	0.0400	1.0
0.05	0.10	0.0500	0.0500	1.0
0.06	0.12	0.0600	0.0600	1.0
0.07	0.14	0.0700	0.0700	1.0
0.08	0.16	0.0800	0.0800	1.0
0.09	0.18	0.0900	0.0900	1.0
0.10	0.20	0.1000	0.1000	1.0

TABLE 11. *Estimated values of β_{max} and $\beta_{\text{max}}/\beta_{\text{min}}$ for various values of β_{min} and β_{max}*

β_{min}	β_{max}	β_{max}	β_{max}	β_{max}
0.01	0.02	0.0100	0.0100	1.0
0.02	0.04	0.0200	0.0200	1.0
0.03	0.06	0.0300	0.0300	1.0
0.04	0.08	0.0400	0.0400	1.0
0.05	0.10	0.0500	0.0500	1.0
0.06	0.12	0.0600	0.0600	1.0
0.07	0.14	0.0700	0.0700	1.0
0.08	0.16	0.0800	0.0800	1.0
0.09	0.18	0.0900	0.0900	1.0
0.10	0.20	0.1000	0.1000	1.0

Table XCV.

HYDRATES.

m.	α .	L.	Δ/m .	L'.	M.	H.
0.2	.541	4.88	6.28	6.25	12.2	61.0
0.4	.430	4.26	6.79	6.72	20.3	50.7
0.6	.352	3.82	7.55	7.41	26.9	44.8
0.8	.292	3.45	8.33	8.12	31.7	39.6
1.0	.241	3.21	9.15	8.85	35.4	35.4
1.2	.201	2.98	10.09	9.70	38.5	32.1
1.3617	.173	2.82	10.22	10.23	40.2	29.5

MIXTURE OF FERRIC CHLORIDE AND ALUMINIUM CHLORIDE.

Table XCVI. FREEZING-POINT MEASUREMENTS.

m.	P.	Δ .
0.4 FeCl ₃	$\frac{1}{1}$	
0.4 AlCl ₃	"	29.38
0.8 FeCl ₃	$\frac{1}{1}$	
0.8 AlCl ₃	"	7.238
1.2 FeCl ₃	$\frac{1}{1}$	
1.2 AlCl ₃	"	13.69
1.6 FeCl ₃	$\frac{1}{1}$	
1.6 AlCl ₃	"	22.5
2.0 FeCl ₃	$\frac{1}{1}$	
2.0 AlCl ₃	"	36.0
2.4 FeCl ₃	$\frac{1}{1}$	
2.4 AlCl ₃	"	56.0
2.7334 FeCl ₃	$\frac{1}{1}$	
2.7334 AlCl ₃	"	84.6

It was necessary to extrapolate this last value, 84.6, since the solution failed to freeze in the freezing mixture.

COLUMBIAN					THE RANGE	
1911	1912	1913	1914	1915	1916	1917
10.00	10.00	10.00	10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00	10.00	10.00	10.00

THE RANGE OF THE COLUMBIAN IS THE RANGE OF THE COLUMBIAN

COLUMBIAN		THE RANGE	
1911	1912	1913	1914
10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00
10.00	10.00	10.00	10.00

THE RANGE OF THE COLUMBIAN IS THE RANGE OF THE COLUMBIAN

THE RANGE OF THE COLUMBIAN IS THE RANGE OF THE COLUMBIAN

employed, i.e., solid carbon dioxide and alcohol. The more concentrated solutions became so viscous, when cooled down, that it was difficult to stir them properly.

Table XCVII. WEIGHT-NORMAL CORRECTIONS.

m.	P.	$W_{\text{Sol.}}$	$W_{\text{Salt.}}$	$W_{\text{H}_2\text{O.}}$	Correction Per Cent.
0.4 FeCl_3	$\frac{1}{11}$	26.2807	1.4786	24.8021	0.8
0.4 AlCl_3	"				
0.8 FeCl_3	$\frac{1}{11}$	27.4565	2.9571	24.4794	2.1
0.8 AlCl_3	"				
1.2 FeCl_3	$\frac{1}{11}$	28.5660	4.4356	24.1504	3.5
1.2 AlCl_3	"				
1.6 FeCl_3	$\frac{1}{11}$	29.7105	5.9142	23.7963	4.8
1.6 AlCl_3	"				
2.0 FeCl_3	$\frac{1}{11}$	30.8182	7.3925	23.4257	6.3
2.0 AlCl_3	"				
2.4 FeCl_3	$\frac{1}{11}$	31.9182	8.8713	23.0469	7.8
2.4 AlCl_3	"				
2.7234 FeCl_3	$\frac{1}{11}$	32.7777	10.0667	22.7110	9.2
2.7254 AlCl_3	"				

It is to be noted that the conductivity measurements, which are given in the following table, are the nearest approximations, as measurements of dissociation. The conductivity measurements were made at zero, whereas some of these mixtures freeze very considerably below 0°C . There is marked hydrolysis, in some cases, as already mentioned, leading to a precipitation of the iron. There is considerable difference between the viscosity of the single solutions

and of the mixtures where the normalities are the same. The assumption that these salts dissociate at once, in the most concentrated solutions, into $\frac{1}{2}\overline{M}, \overline{Cl}, \overline{Cl}$, is certainly not correct. Finally, the values of Λ_{∞} are calculated values and may differ considerably from the true values, because of the factors just mentioned.

THESE ARE THE RESULTS OF THE WORK DONE BY THE
 THE RESEARCHERS IN THE FIELD OF THE
 THE RESEARCHERS IN THE FIELD OF THE
 THE RESEARCHERS IN THE FIELD OF THE
 THE RESEARCHERS IN THE FIELD OF THE
 THE RESEARCHERS IN THE FIELD OF THE

Table XCVIII.

CONDUCTIVITY MEASURED AT 25°.

M.	P.	Γ_0 .	V_c .	k_w .	k.	D.	k_e .	Λ_v	α .
0.4 FeCl ₃	1	0.2	5.000	.040318	.02780	.007465	.01955	97.6	0.140
0.4 AlCl ₃	"	"	"		.027937		.020568	101.8	.577
0.8 Fe	1	0.4	2.500	.060462	.03786	.017928	.026928	67.3	.700
0.8 Al	"	"	"		.04053		.033534	83.7	.461
1.2 Fe	1	0.6	1.6666	.055374	.04645	.043691	.019059	31.8	.144
1.2 Al	"	"	"		.058115		.036815	61.4	.532
1.6 Fe	1	0.8	1.250	.056904	.05147	.04575	.01604	20.1	.091
1.6 Al	"	"	"		.06118		.04086	51.1	.321
2.0 Fe	1	1.0	1.00	.047916	.053147	.071836	.006597	6.6	.070
2.0 Al	"	"	"		.066595		.041319	41.3	.227
2.4 Fe	1	1.2	0.8333	.037666	.053145	.083699	.0000	00.0	.00
2.4 Al	"	"	"		.06842		.03787	31.6	.174
2.7334 Fe	1	1.26170	.7344	.039595	.051777	.089762	.0000	00.0	.00
" Al	"	"	"		.06760		.029595	21.7	.120

Table XCIX.

HYDRATES.

M.	P.	n_D .	α .	L_c .	M.	Δ .	Spdr. Cor.	M_c .	H_c .
0.4 Fe Cl ₃	1	0.2	0.440	4.31	12.2	29.78	0.8	13.5	27.5
0.4 AlCl ₃	"	"	.577	5.08	7.4			8.2	41.0
0.8 Fe	1	0.4	.506	3.57	20.3	7.238	2.1	14.6	49.0
0.8 Al	"	"	.461	4.44	17.4			10.6	41.5
1.2 Fe	1	0.6	.144	3.67	26.9	13.69	3.5	25.8	47.0
1.2 Al	"	"	.338	5.74	25.5			21.4	40.7
1.6 Fe	1	0.8	.091	2.37	31.7	22.5	4.8	29.0	76.2
1.6 Al	"	"	.231	5.43	31.6			28.9	70.1
2.0 Fe	1	1.0	.050	2.03	35.4	36.0	6.3	32.2	32.2
2.0 Al	"	"	.227	5.02	36.7			33.7	73.7
2.4 Fe	1	1.2	.00	1.86	38.5	56.0	7.8	74.6	28.8
2.4 Al	"	"	.174	2.83	40.4			36.7	30.6
2.72 ⁸⁴ Fe	1	1.36	17.00	1.86	40.2	84.6	9.2	36.1	26.5
" Al	"	"	.120	2.53	43.4			39.6	39.0

200 200 200 200 200 200 200 200

200 200 200 200 200 200 200 200

200 200 200 200 200 200 200 200

200 200 200 200 200 200 200 200

200 200 200 200 200 200 200 200

200 200 200 200 200 200 200 200

200 200 200 200 200 200 200 200

200 200 200 200 200 200 200 200

200 200 200 200 200 200 200 200

200 200 200 200 200 200 200 200

These values of M and of M_0 are plotted in curves in Figure XVI. The general relations shown by these curves are the same as have previously been pointed out. The fact that the hydrating power of both the aluminium chloride and the ferric chloride is apparently greater in the mixture than in the separate solutions, in the most dilute solutions employed, is probably due to the effect of hydrolysis in increasing the value of L , and consequently diminishing M , though possibly partially due to change in dissociation and difference in the hydrating power of the ions and molecules. It should be noted, also, that the hydrating power of the aluminium chloride appears to be relatively less than that of the ferric chloride in the more dilute solutions; but that this relation is reversed in the more concentrated solutions. This may be due to hydrolysis, since if the aluminium chloride hydrolyzes more rapidly than the ferric chloride, then, in the more dilute solutions, where the effects of hydrolysis would be pronounced, the result would be to increase the value of L and, consequently, to diminish the value of M , based upon it.

In Tables C and CI the abbreviations have the usual significance.

Table C. ALUMINIUM CHLORIDE IN MIXTURE OF ALUMINIUM CHLORIDE AND FERRIC CHLORIDE.

m.	D_{H_2O} .	D_M .
0.2	246	-0.8
0.4	367	0.8
0.6	488	1.1
0.8	552	2.7
1.0	618	3.0
1.2	671	3.7
1.3617	709	3.8

Table CI. FERRIC CHLORIDE IN MIXTURE OF ALUMINIUM CHLORIDE AND FERRIC CHLORIDE.

m.	D_{H_2O} .	D_M .
0.2	150	-1.3
0.4	308	0.7
0.6	456	1.1
0.8	543	2.7
1.0	636	3.2
1.2	699	3.9
1.3617	758	4.1

These values are plotted in curves in Figures XVII and XVIII.

TABLE 1. Values of the function $f(x)$ for x from 0 to 1.0 in increments of 0.1.

TABLE 2. Values of the function $f(x)$ for x from 0 to 1.0 in increments of 0.05.

TABLE 3. Values of the function $f(x)$ for x from 0 to 1.0 in increments of 0.02.

x	$f(x)$	$f'(x)$
0.0	0.0000	0.0000
0.1	0.0950	0.9050
0.2	0.3600	0.6400
0.3	0.6750	0.3250
0.4	0.9200	0.0800
0.5	0.9900	0.0100
0.6	0.9950	0.0050
0.7	0.9975	0.0025
0.8	0.9988	0.0012
0.9	0.9995	0.0005
1.0	1.0000	0.0000

TABLE 4. Values of the function $f(x)$ for x from 0 to 1.0 in increments of 0.01.

TABLE 5. Values of the function $f(x)$ for x from 0 to 1.0 in increments of 0.005.

x	$f(x)$	$f'(x)$
0.0	0.0000	0.0000
0.05	0.0475	0.9525
0.1	0.0950	0.9050
0.15	0.1425	0.8575
0.2	0.1900	0.8100
0.25	0.2375	0.7625
0.3	0.2850	0.7150
0.35	0.3325	0.6675
0.4	0.3800	0.6200
0.45	0.4275	0.5725
0.5	0.4750	0.5250
0.55	0.5225	0.4775
0.6	0.5700	0.4300
0.65	0.6175	0.3825
0.7	0.6650	0.3350
0.75	0.7125	0.2875
0.8	0.7600	0.2400
0.85	0.8075	0.1925
0.9	0.8550	0.1450
0.95	0.9025	0.0975
1.0	0.9500	0.0500


These values are plotted in Figure 1 to show the shape of the function.

Figure 1.

OTHER FACTORS INFLUENCING THE AMOUNT OF WATER REQUIRED AS SOLVENT IN THE SINGLE SOLUTION OF ALUMINUM CHLORIDE AND IN THE MIXTURE OF PERBIC CHLORIDE AND ALUMINUM CHLORIDE. THE VALUES OF λ FOR ALUMINUM CHLORIDE IN SINGLE SOLUTION AND IN THE MIXTURE OF ALUMINUM CHLORIDE AND PERBIC CHLORIDE.

DEPT. EXPERIMENTAL ENGINEERING, SIKLEY COLLEGE, CORNELL UNIVERSITY.

I. GRAMS OF WATER.




GRAM MOLECULES OF SALT PER LITER OF SOLUTION.

FIGURE XVII.

DIFFERENCE BETWEEN THE VALUES OF THE PERKINS CHLORIDE IN PURE SOLUTION AND IN THE MIXTURE OF PERKINS CHLORIDE AND ALUMINUM CHLORIDE.

DIFFERENCE BETWEEN THE VALUES OF THE PERKINS CHLORIDE IN PURE SOLUTION AND IN THE MIXTURE OF PERKINS CHLORIDE AND ALUMINUM CHLORIDE.



The effect of decrease in the amount of water present as solvent is plainly shown in the diminishing values of Δ as the amount of solvent is lessened. The slight irregularities are probably due to unavoidable experimental error, and inaccuracy of the conductivity method as a measure of dissociation.

Thus far all of the Salts Studied have a Common Anion. In order to Extend the Investigation to a Pair of Salts having a Common Cation, a Mixture of Calcium Nitrate and Calcium chloride was employed.

The data for the separate solutions of these two salts has already been given. The results obtained with the mixture follow.

Table CII. FREEZING-POINT MEASUREMENTS.

m.	P.	Δ .
0.5 Ca_2NO_3	1	
0.5 CaCl_2	"	2.50
0.9 Ca_2NO_3	1	
0.9 CaCl_2	"	4.88
1.3 Ca_2NO_3	1	
1.3 CaCl_2	"	7.67
1.7 Ca_2NO_3	1	
1.7 CaCl_2	"	11.00
2.1 Ca_2NO_3	1	
2.1 CaCl_2	"	14.78
2.5 Ca_2NO_3	1	
2.5 CaCl_2	"	19.2
2.9 Ca_2NO_3	1	
2.9 CaCl_2	"	24.2
3.31102 Ca_2NO_3	1	
3.31102 CaCl_2	"	30.7

Table CIII.
CONDUCTIVITY MEASUREMENTS.

m.	P.	m _c .	v _c .	k _v	k.	D.	k _c .	Λ_v .	α .
0.5 Ca(NO ₃) ₂	1 "	0.25 "	4.0000	.039475	.020975	.05573	.018254	73.0	0.547
0.5 CaCl ₂			"		.024073		.021231	84.9	.575
0.9 Ca(NO ₃) ₂	1 "	0.45 "	2.2222	.059937	.03259	.018010	.026780	59.5	.42
0.9 CaCl ₂			"		.039688		.073157	73.7	.199
1.3 Ca(NO ₃) ₂	1 "	0.65 "	1.538	.073550	.042932	.023618	.030233	47.4	.353
1.7 CaCl ₂			"		.054236		.042716	65.7	.445
1.7 Ca(NO ₃) ₂	1 "	0.85 "	1.1765	.081797	.050272	.035578	.031691	37.3	.277
1.7 CaCl ₂			"		.066903		.050106	59.0	.400
2.1 Ca(NO ₃) ₂	1 "	1.05 "	0.9523	.08410	.056501	.050445	.029266	27.9	.207
2.1 CaCl ₂			"		.072344		.054654	52.2	.754
2.5 Ca(NO ₃) ₂	1 "	1.25 "	0.8000	.082681	.060363	.066555	.023751	19.0	.141
2.5 CaCl ₂			"		.088853		.058930	47.1	.320
2.9 Ca(NO ₃) ₂	1 "	1.45 "	.68966	.077154	.063236	.053467	.016375	11.8	.084
2.9 CaCl ₂			"		.097385		.060780	41.	.284
3.11 Ca(NO ₃) ₂	1 "	1.6555 "	.60405	.069855	.064674	.068121	.008650	2.2	.076
" CaCl ₂			"		.00330		.061202	37.0	.251

Year	Month	Day	Time	Location	Activity	Remarks
1911	Jan	1	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	2	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	3	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	4	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	5	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	6	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	7	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	8	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	9	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	10	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	11	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	12	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	13	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	14	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	15	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	16	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	17	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	18	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	19	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	20	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	21	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	22	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	23	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	24	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	25	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	26	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	27	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	28	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	29	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	30	10:00	St. Paul	Prayer	Prayer meeting
1911	Jan	31	10:00	St. Paul	Prayer	Prayer meeting

Table CIV. WEIGHT-NORMAL CORRECTIONS.

m.	P.	W _{Sol.}	W _{Salt.}	W _{H₂O.}	Correction Per Cent.
0.5 Ca ₂ NO ₃	1	26.3336	1.7197	24.6139	1.5
0.5 CaCl ₂	"				
0.9 Ca ₂ NO ₃	1	27.3415	3.0954	24.2461	3.0
0.9 CaCl ₂	"				
1.3 Ca ₂ NO ₃	1	28.3615	4.4711	23.8904	4.4
1.3 CaCl ₂	"				
1.7 Ca ₂ NO ₃	1	29.3082	5.8469	23.4613	6.2
1.7 CaCl ₂	"				
2.1 Ca ₂ NO ₃	1	30.2933	7.326	23.0707	7.7
2.1 CaCl ₂	"				
2.5 Ca ₂ NO ₃	1	31.2197	8.5984	22.6213	9.5
2.5 CaCl ₂	"				
2.9 Ca ₂ NO ₃	1	32.1762	9.9741	22.2021	11.2
2.9 CaCl ₂	"				
3.3111 Ca ₂ NO ₃	1	33.0792	11.3878	21.6914	13.2
3.311 CaCl ₂	"				

Mathematical Summary

Order	Class	Subclass	Group	Count	Mean
1.1	1.1.1	1.1.1.1	1.1.1.1.1	1	1.1.1.1.1
1.2	1.2.1	1.2.1.1	1.2.1.1.1	1	1.2.1.1.1
1.3	1.3.1	1.3.1.1	1.3.1.1.1	1	1.3.1.1.1
1.4	1.4.1	1.4.1.1	1.4.1.1.1	1	1.4.1.1.1
1.5	1.5.1	1.5.1.1	1.5.1.1.1	1	1.5.1.1.1
1.6	1.6.1	1.6.1.1	1.6.1.1.1	1	1.6.1.1.1
1.7	1.7.1	1.7.1.1	1.7.1.1.1	1	1.7.1.1.1
1.8	1.8.1	1.8.1.1	1.8.1.1.1	1	1.8.1.1.1
1.9	1.9.1	1.9.1.1	1.9.1.1.1	1	1.9.1.1.1
1.10	1.10.1	1.10.1.1	1.10.1.1.1	1	1.10.1.1.1
1.11	1.11.1	1.11.1.1	1.11.1.1.1	1	1.11.1.1.1
1.12	1.12.1	1.12.1.1	1.12.1.1.1	1	1.12.1.1.1
1.13	1.13.1	1.13.1.1	1.13.1.1.1	1	1.13.1.1.1
1.14	1.14.1	1.14.1.1	1.14.1.1.1	1	1.14.1.1.1
1.15	1.15.1	1.15.1.1	1.15.1.1.1	1	1.15.1.1.1
1.16	1.16.1	1.16.1.1	1.16.1.1.1	1	1.16.1.1.1
1.17	1.17.1	1.17.1.1	1.17.1.1.1	1	1.17.1.1.1
1.18	1.18.1	1.18.1.1	1.18.1.1.1	1	1.18.1.1.1
1.19	1.19.1	1.19.1.1	1.19.1.1.1	1	1.19.1.1.1
1.20	1.20.1	1.20.1.1	1.20.1.1.1	1	1.20.1.1.1
1.21	1.21.1	1.21.1.1	1.21.1.1.1	1	1.21.1.1.1
1.22	1.22.1	1.22.1.1	1.22.1.1.1	1	1.22.1.1.1
1.23	1.23.1	1.23.1.1	1.23.1.1.1	1	1.23.1.1.1
1.24	1.24.1	1.24.1.1	1.24.1.1.1	1	1.24.1.1.1
1.25	1.25.1	1.25.1.1	1.25.1.1.1	1	1.25.1.1.1
1.26	1.26.1	1.26.1.1	1.26.1.1.1	1	1.26.1.1.1
1.27	1.27.1	1.27.1.1	1.27.1.1.1	1	1.27.1.1.1
1.28	1.28.1	1.28.1.1	1.28.1.1.1	1	1.28.1.1.1
1.29	1.29.1	1.29.1.1	1.29.1.1.1	1	1.29.1.1.1
1.30	1.30.1	1.30.1.1	1.30.1.1.1	1	1.30.1.1.1
1.31	1.31.1	1.31.1.1	1.31.1.1.1	1	1.31.1.1.1
1.32	1.32.1	1.32.1.1	1.32.1.1.1	1	1.32.1.1.1
1.33	1.33.1	1.33.1.1	1.33.1.1.1	1	1.33.1.1.1
1.34	1.34.1	1.34.1.1	1.34.1.1.1	1	1.34.1.1.1
1.35	1.35.1	1.35.1.1	1.35.1.1.1	1	1.35.1.1.1
1.36	1.36.1	1.36.1.1	1.36.1.1.1	1	1.36.1.1.1
1.37	1.37.1	1.37.1.1	1.37.1.1.1	1	1.37.1.1.1
1.38	1.38.1	1.38.1.1	1.38.1.1.1	1	1.38.1.1.1
1.39	1.39.1	1.39.1.1	1.39.1.1.1	1	1.39.1.1.1
1.40	1.40.1	1.40.1.1	1.40.1.1.1	1	1.40.1.1.1
1.41	1.41.1	1.41.1.1	1.41.1.1.1	1	1.41.1.1.1
1.42	1.42.1	1.42.1.1	1.42.1.1.1	1	1.42.1.1.1
1.43	1.43.1	1.43.1.1	1.43.1.1.1	1	1.43.1.1.1
1.44	1.44.1	1.44.1.1	1.44.1.1.1	1	1.44.1.1.1
1.45	1.45.1	1.45.1.1	1.45.1.1.1	1	1.45.1.1.1
1.46	1.46.1	1.46.1.1	1.46.1.1.1	1	1.46.1.1.1
1.47	1.47.1	1.47.1.1	1.47.1.1.1	1	1.47.1.1.1
1.48	1.48.1	1.48.1.1	1.48.1.1.1	1	1.48.1.1.1
1.49	1.49.1	1.49.1.1	1.49.1.1.1	1	1.49.1.1.1
1.50	1.50.1	1.50.1.1	1.50.1.1.1	1	1.50.1.1.1
1.51	1.51.1	1.51.1.1	1.51.1.1.1	1	1.51.1.1.1
1.52	1.52.1	1.52.1.1	1.52.1.1.1	1	1.52.1.1.1
1.53	1.53.1	1.53.1.1	1.53.1.1.1	1	1.53.1.1.1
1.54	1.54.1	1.54.1.1	1.54.1.1.1	1	1.54.1.1.1
1.55	1.55.1	1.55.1.1	1.55.1.1.1	1	1.55.1.1.1
1.56	1.56.1	1.56.1.1	1.56.1.1.1	1	1.56.1.1.1
1.57	1.57.1	1.57.1.1	1.57.1.1.1	1	1.57.1.1.1
1.58	1.58.1	1.58.1.1	1.58.1.1.1	1	1.58.1.1.1
1.59	1.59.1	1.59.1.1	1.59.1.1.1	1	1.59.1.1.1
1.60	1.60.1	1.60.1.1	1.60.1.1.1	1	1.60.1.1.1
1.61	1.61.1	1.61.1.1	1.61.1.1.1	1	1.61.1.1.1
1.62	1.62.1	1.62.1.1	1.62.1.1.1	1	1.62.1.1.1
1.63	1.63.1	1.63.1.1	1.63.1.1.1	1	1.63.1.1.1
1.64	1.64.1	1.64.1.1	1.64.1.1.1	1	1.64.1.1.1
1.65	1.65.1	1.65.1.1	1.65.1.1.1	1	1.65.1.1.1
1.66	1.66.1	1.66.1.1	1.66.1.1.1	1	1.66.1.1.1
1.67	1.67.1	1.67.1.1	1.67.1.1.1	1	1.67.1.1.1
1.68	1.68.1	1.68.1.1	1.68.1.1.1	1	1.68.1.1.1
1.69	1.69.1	1.69.1.1	1.69.1.1.1	1	1.69.1.1.1
1.70	1.70.1	1.70.1.1	1.70.1.1.1	1	1.70.1.1.1
1.71	1.71.1	1.71.1.1	1.71.1.1.1	1	1.71.1.1.1
1.72	1.72.1	1.72.1.1	1.72.1.1.1	1	1.72.1.1.1
1.73	1.73.1	1.73.1.1	1.73.1.1.1	1	1.73.1.1.1
1.74	1.74.1	1.74.1.1	1.74.1.1.1	1	1.74.1.1.1
1.75	1.75.1	1.75.1.1	1.75.1.1.1	1	1.75.1.1.1
1.76	1.76.1	1.76.1.1	1.76.1.1.1	1	1.76.1.1.1
1.77	1.77.1	1.77.1.1	1.77.1.1.1	1	1.77.1.1.1
1.78	1.78.1	1.78.1.1	1.78.1.1.1	1	1.78.1.1.1
1.79	1.79.1	1.79.1.1	1.79.1.1.1	1	1.79.1.1.1
1.80	1.80.1	1.80.1.1	1.80.1.1.1	1	1.80.1.1.1
1.81	1.81.1	1.81.1.1	1.81.1.1.1	1	1.81.1.1.1
1.82	1.82.1	1.82.1.1	1.82.1.1.1	1	1.82.1.1.1
1.83	1.83.1	1.83.1.1	1.83.1.1.1	1	1.83.1.1.1
1.84	1.84.1	1.84.1.1	1.84.1.1.1	1	1.84.1.1.1
1.85	1.85.1	1.85.1.1	1.85.1.1.1	1	1.85.1.1.1
1.86	1.86.1	1.86.1.1	1.86.1.1.1	1	1.86.1.1.1
1.87	1.87.1	1.87.1.1	1.87.1.1.1	1	1.87.1.1.1
1.88	1.88.1	1.88.1.1	1.88.1.1.1	1	1.88.1.1.1
1.89	1.89.1	1.89.1.1	1.89.1.1.1	1	1.89.1.1.1
1.90	1.90.1	1.90.1.1	1.90.1.1.1	1	1.90.1.1.1
1.91	1.91.1	1.91.1.1	1.91.1.1.1	1	1.91.1.1.1
1.92	1.92.1	1.92.1.1	1.92.1.1.1	1	1.92.1.1.1
1.93	1.93.1	1.93.1.1	1.93.1.1.1	1	1.93.1.1.1
1.94	1.94.1	1.94.1.1	1.94.1.1.1	1	1.94.1.1.1
1.95	1.95.1	1.95.1.1	1.95.1.1.1	1	1.95.1.1.1
1.96	1.96.1	1.96.1.1	1.96.1.1.1	1	1.96.1.1.1
1.97	1.97.1	1.97.1.1	1.97.1.1.1	1	1.97.1.1.1
1.98	1.98.1	1.98.1.1	1.98.1.1.1	1	1.98.1.1.1
1.99	1.99.1	1.99.1.1	1.99.1.1.1	1	1.99.1.1.1
2.00	2.00.1	2.00.1.1	2.00.1.1.1	1	2.00.1.1.1
2.01	2.01.1	2.01.1.1	2.01.1.1.1	1	2.01.1.1.1
2.02	2.02.1	2.02.1.1	2.02.1.1.1	1	2.02.1.1.1
2.03	2.03.1	2.03.1.1	2.03.1.1.1	1	2.03.1.1.1
2.04	2.04.1	2.04.1.1	2.04.1.1.1	1	2.04.1.1.1
2.05	2.05.1	2.05.1.1	2.05.1.1.1	1	2.05.1.1.1
2.06	2.06.1	2.06.1.1	2.06.1.1.1	1	2.06.1.1.1
2.07	2.07.1	2.07.1.1	2.07.1.1.1	1	2.07.1.1.1
2.08	2.08.1	2.08.1.1	2.08.1.1.1	1	2.08.1.1.1
2.09	2.09.1	2.09.1.1	2.09.1.1.1	1	2.09.1.1.1
2.10	2.10.1	2.10.1.1	2.10.1.1.1	1	2.10.1.1.1
2.11	2.11.1	2.11.1.1	2.11.1.1.1	1	2.11.1.1.1
2.12	2.12.1	2.12.1.1	2.12.1.1.1	1	2.12.1.1.1
2.13	2.13.1	2.13.1.1	2.13.1.1.1	1	2.13.1.1.1
2.14	2.14.1	2.14.1.1	2.14.1.1.1	1	2.14.1.1.1
2.15	2.15.1	2.15.1.1	2.15.1.1.1	1	2.15.1.1.1
2.16	2.16.1	2.16.1.1	2.16.1.1.1	1	2.16.1.1.1
2.17	2.17.1	2.17.1.1	2.17.1.1.1	1	2.17.1.1.1
2.18	2.18.1	2.18.1.1	2.18.1.1.1	1	2.18.1.1.1
2.19	2.19.1	2.19.1.1	2.19.1.1.1	1	2.19.1.1.1
2.20	2.20.1	2.20.1.1	2.20.1.1.1	1	2.20.1.1.1
2.21	2.21.1	2.21.1.1	2.21.1.1.1	1	2.21.1.1.1
2.22	2.22.1	2.22.1.1	2.22.1.1.1	1	2.22.1.1.1
2.23	2.23.1	2.23.1.1	2.23.1.1.1	1	2.23.1.1.1
2.24	2.24.1	2.24.1.1	2.24.1.1.1	1	2.24.1.1.1
2.25	2.25.1	2.25.1.1	2.25.1.1.1	1	2.25.1.1.1
2.26	2.26.1	2.26.1.1	2.26.1.1.1	1	2.26.1.1.1
2.27	2.27.1	2.27.1.1	2.27.1.1.1	1	2.27.1.1.1
2.28	2.28.1	2.28.1.1	2.28.1.1.1	1	2.28.1.1.1
2.29	2.29.1	2.29.1.1	2.29.1.1.1	1	2.29.1.1.1
2.30	2.30.1	2.30.1.1	2.30.1.1.1	1	2.30.1.1.1
2.31	2.31.1	2.31.1.1	2.31.1.1.1	1	2.31.1.1.1
2.32	2.32.1	2.32.1.1	2.32.1.1.1	1	2.32.1.1.1
2.33	2.33.1	2.33.1.1	2.33.1.1.1	1	2.33.1.1.1
2.34	2.34.1	2.34.1.1	2.34.1.1.1	1	2.34.1.1.1
2.35	2.35.1	2.35.1.1	2.35.1.1.1	1	2.35.1.1.1
2.36	2.36.1	2.36.1.1	2.36.1.1.1	1	2.36.1.1.1
2.37	2.37.1	2.37.1.1	2.37.1.1.1	1	2.37.1.1.1
2.38	2.38.1	2.38.1.1	2.38.1.1.1	1	2.38.1.1.1
2.39	2.39.1	2.39.1.1	2.39.1.1.1	1	2.39.1.1.1
2.40	2.40.1	2.40.1.1	2.40.1.1.1	1	2.40.1.1.1
2.41	2.41.1	2.41.1.1	2.41.1.1.1	1	2.41.1.1.1
2.42	2.42.1	2.42.1.1	2.42.1.1.1	1	2.42.1.1.1
2.43	2.43.1	2.43.1.1	2.43.1.1.1	1	2.43.1.1.1
2.44	2.44.1	2.44.1.1	2.44.1.1.1	1	2.44.1.1.1
2.45	2.45.1	2.45.1.1	2.45.1.1.1	1	2.45.1.1.1
2.46	2.46.1	2.46.1.1	2.46.1.1.1	1	2.46.1.1.1
2.47	2.47.1	2.47.1.1	2.47.1.1.1	1	2.47.1.1.1
2.48	2.48.1	2.48.1.1	2.48.1.1.1	1	2.48.1.1.1
2.49	2.49.1	2.49.1.1	2.49.1.1.1	1	2.49.1.1.1
2.50	2.50.1	2.50.1.1	2.50.1.1.1	1	2.50.1.1.1
2.51	2.51.1	2.51.1.1	2.51.1.1.1	1	2.51.1.1.1
2.52	2.52.1	2.52.1.1	2.52.1.1.1	1	2.52.1.1.1
2.53	2.53.1	2.53.1.1	2.53.1.1.1	1	2.53.1.1.1
2.54	2.54.1	2.54.1.1	2.54.1.1.1	1	2.54.1.1.1
2.55	2.55.1	2.55.1.1	2.55.1.1.1	1	2.55.1.1.1
2.56	2.56.1	2.56.1.1	2.56.1.1.1	1	2.56.1.1.1
2.57	2.57.1	2.57.1.1	2.57.1.1.1	1	2.57.1.1.1
2.58	2.58.1	2.58.1.1	2.58.1.1.1	1	2.58.1.1.1
2.59	2.59.1	2.59.1.1	2.59.1.1.1	1	2.59.1.1.1
2.60	2.60.1	2.60.1.1	2.60.1.1.1	1	2.60.1.1.1
2.61	2.61.1	2.61.1.1	2.61.1.1.1	1	2.61.1.1.1
2.62	2.62.1	2.62.1.1	2.62.1.1.1	1	2.62.1.1.1
2.63	2.63.1	2.63.1.1	2.63.1.1.1	1	2.63.1.1.1
2.64	2.64.1	2.64.1.1	2.64.1		

HYDRATES.

Table CV.

n.	P.	m _c .	α.	L _c .	M.	Δ.	WT. Normal Cor.	M _c .	H _c .
0.5 Ca2NO ₃	1	0.25	0.543	3.88	6.5	2.50	1.5	4.6	18.4
0.5 CaCl ₂	"	"	.575	4.00	8.2			7.0	28.0
0.9 Ca2NO ₃	1	0.45	.442	3.51	8.0	4.88	3.0	7.4	16.4
0.9 CaCl ₂	"	"	.499	3.72	11.7			11.2	24.9
1.3 Ca2NO ₃	1	0.65	.353	3.17	10.5	7.67	4.4	9.5	14.8
1.3 CaCl ₂	"	"	.445	3.52	15.7			15.1	23.2
1.7 Ca2NO ₃	1	0.85	.277	3.89	13.1	11.00	6.2	13.1	14.2
1.7 CaCl ₂	"	"	.400	3.35	18.9			18.2	21.4
2.1 Ca2NO ₃	1	1.05	.207	2.63	15.7	14.78	7.7	14.1	13.4
2.1 CaCl ₂	"	"	.354	3.18	21.9			20.6	19.8
2.5 Ca2NO ₃	1	1.25	.141	2.39	18.4	19.20	9.5	16.4	13.1
2.5 CaCl ₂	"	"	.320	3.05	24.3			22.8	18.2
2.9 Ca2NO ₃	1	1.45	.084	2.17	20.4	24.2	11.2	17.6	12.1
2.9 CaCl ₂	"	"	.284	2.92	27.3			25.2	17.4
3.311 Ca2NO ₃	1	1.655b	.039	2.01	22.4	50.7	13.2	16.4	9.9
3.311 CaCl ₂	"	"	.251	2.79	30.1			25.6	15.5

CASHEW OILSEED, 4.1%.


" " MIXED WITH CALCIUM NITRATE.

" NITRATE ALONE.

" " MIXED WITH CALCIUM CHLORIDE.

DEPT. EXPERIMENTAL ENGINEERING, ARLY COLLEGE, CORNELL UNIVERSITY.

NUMBER OF GRAM MOLECULES OF WATER COMBINED.



GRAM MOLECULES OF SALT PER LITER OF SOLUTION.

FIGURE XIX.

The above values of M are plotted in curves in Figure XIX. The results are of the same character as those shown by the preceding mixtures. The most concentrated solution measured shows a somewhat lower value for M in the mixture than we should expect, but it must be borne in mind that at this great concentration the experimental errors involved are necessarily very large.

Tables CVI and CVII contain the data necessary to compare the change in the amount of solvent present with the change in the values of M in separate solution and in the mixture.

Table CVI. CALCIUM CHLORIDE IN MIXTURE OF CALCIUM NITRATE AND CALCIUM CHLORIDE.

<u>m.</u>	<u>D_{H₂O}.</u>	<u>D_M.</u>
0.25	92	1.2
0.45	153	0.5
0.65	202	0.6
0.85	260	0.7
1.05	306	1.1
1.25	358	1.5
1.45	392	2.1
1.6555	386	4.5

Table CVII. CALCIUM NITRATE IN MIXTURE OF CALCIUM NITRATE
AND CALCIUM CHLORIDE.

<u>m.</u>	<u>D_{H₂O.}</u>	<u>D_{L.}</u>
0.35	181	1.7
0.45	313	0.6
0.65	388	0.9
0.85	350	1.0
1.05	404	1.6
1.25	447	2.0
1.45	496	2.8
1.6555	575	6.0

These values are plotted in curves in Figures XX and XXI. A diminution in the amount of water present as solvent causes a diminution in the value of M. This effect appears to be greater with increasing concentration. It is probable that in the most dilute solutions the difference in the hydrating power of the ions and molecules affects the curve, changing its direction.


TABLE 1. MEAN VALUES OF THE DIFFERENT PARAMETERS OF THE STUDY

PARAMETER	MEAN	STANDARD DEVIATION
1. Age	25.5	2.5
2. Height	1.75	0.05
3. Weight	70.0	10.0
4. BMI	22.5	2.5
5. Systolic BP	120	10
6. Diastolic BP	80	10
7. Heart rate	72	10
8. Fasting glucose	90	10
9. Fasting insulin	10	5

The mean values of the different parameters of the study are presented in Table 1. The mean age of the subjects was 25.5 years, with a standard deviation of 2.5 years. The mean height was 1.75 m, with a standard deviation of 0.05 m. The mean weight was 70.0 kg, with a standard deviation of 10.0 kg. The mean BMI was 22.5, with a standard deviation of 2.5. The mean systolic BP was 120 mmHg, with a standard deviation of 10 mmHg. The mean diastolic BP was 80 mmHg, with a standard deviation of 10 mmHg. The mean heart rate was 72 bpm, with a standard deviation of 10 bpm. The mean fasting glucose was 90 mg/dL, with a standard deviation of 10 mg/dL. The mean fasting insulin was 10 µU/mL, with a standard deviation of 5 µU/mL.

DETERMINED BY THE METHOD OF MIXING AS SOLVENT TOWARD THE CALCIUM ION-
 RATE TO THE MIXING OF CALCIUM NITRATE AND CALCIUM CHLORIDE AND BY THE SEPARATE
 SOLUTION OF CALCIUM NITRATE.
 DETERMINED FROM THE VALUES OF γ FOR CALCIUM NITRATE IN SEPARATE SOLUTION AND
 IN THE MIXTURE OF CALCIUM NITRATE AND CALCIUM CHLORIDE.


DEPT. EXPERIMENTAL ENGINEERING, SIKLEY COLLEGE, CORNELL UNIVERSITY.



II. GRAM MOLECULES OF WATER.

GRAM MOLECULES OF SALT PER LITER OF SOLUTION.
 FIGURE XX.

DETERMINE THE AMOUNT OF WATER REQUIRED TO SOLVATE THE CALCIUM CHLORIDE IN THE MIXTURE OF CALCIUM NITRATE AND CALCIUM CHLORIDE AND IN THE SEPARATE SOLUTION OF CALCIUM CHLORIDE.
 DETERMINE THE VALUES OF η FOR CALCIUM CHLORIDE IN SEPARATE SOLUTION AND IN THE MIXTURE OF CALCIUM NITRATE AND CALCIUM CHLORIDE.



GRAM MOLECULES OF SALT PER LITER OF SOLUTION.
 FIGURE XVI.

THE EFFECT OF CHANGE IN TEMPERATURE UPON THE CONDUCTIVITIES OF SEPARATE SOLUTIONS OF ELECTROLYTES AND UPON MIXTURES OF THESE ELECTROLYTES.

In this paper we have repeatedly called attention to the fact that the conductivity method furnishes us with only an approximate value of the dissociation of a salt. The conductivity method of measuring dissociation is, however, the most general one available. In order to study the effect of change of temperature upon the conductivities of separate solutions of electrolytes, and upon mixtures of these electrolytes it was necessary to select a pair of salts which do not form double salts in solution, and which have little or no hydrating power. Potassium chloride and ammonium chloride were employed.

The conductivities of single solutions of these salts and of their mixtures were studied at 0° and 12°. Comparison having shown that our values agree very closely with those obtained by Jones and West¹ and Jones and Knight², Jones and Knight's values at 25° were used. The values of Jones and West, at 18°, show a very close agreement with those obtained experimentally by Kohlrausch at the same temperature. The

1. Amer. Chem. Journ. 34, 321 (1905)
2. " " " 22, 125 (1899).

values for potassium chloride at 25° are taken from the results of Kohlrausch.

POTASSIUM CHLORIDE.

Table CVIII. MOLECULAR CONDUCTIVITIES.

v.	$\mu_{v0^{\circ}}$.	$\mu_{v12^{\circ}}$.	v.	$\mu_{v25^{\circ}}$.
1	61.58	82.05	1	104.1
2	63.02	84.43	2	108.6
5	65.85	87.84	4	111.2
20	68.59	92.88	16	121.5
100	72.13	98.69	80	128.5
200	73.73	101.3	160	131.1
1000	74.31	103.3	640	132.9

AMMONIUM CHLORIDE.

Table CIX. MOLECULAR CONDUCTIVITIES.

v.	$\mu_{v0^{\circ}}$.	$\mu_{v12^{\circ}}$.	v.	$\mu_{v25^{\circ}}$.
1	61.02	81.15	1	99.3
2	62.53	83.57	2	104.9
5	63.98	87.66	4	110.4
20	68.13	93.19	16	120.9
100	71.64	99.94	80	131.8
200	72.89	101.3	160	134.2
1000	74.10	103.7	640	138.1

Since we are employing some of the values of Jones and West, Jones and Knight and Kohlrausch, at 25°, we have tabulated a few values taken from these sources side by side with our own.

POTASSIUM CHLORIDE.

Table CX. COMPARISON OF CONDUCTIVITIES.

v.	Ours.	Jones and West.	(Jones and West.)	Kohlrausch.
	$\mu_{v0^{\circ}}$.	$\mu_{v0^{\circ}}$.	$\mu_{v18^{\circ}}$.	$\mu_{v18^{\circ}}$.
1	61.58	----	---	---
2	63.02	62.96	95.9	95.8
5	65.85	----	---	---

Table CX. COMPARISON OF CONDUCTIVITIES. (Cont.)

v.	$\mu_v 0^\circ$.	$\mu_v 0^\circ$.	$\mu_v 18^\circ$.	$\mu_v 18^\circ$.
8	---	66.47	103.6	----
16	---	68.4	107.0	----
20	68.59	---	----	108.3
32	---	70.27	110.7	----
120	72.13	---	----	114.7
128	---	73.0	115.5	----
200	73.73	---	----	----
1000	74.31	---	----	119.3
1024	---	75.14	119.3	----

AMMONIUM CHLORIDE.

Table CXI. COMPARISON OF CONDUCTIVITIES.

v.	Ours. $\mu_v 0^\circ$.	Jones West. $\mu_v 0^\circ$.	Jones West. $\mu_v 18^\circ$.	Kohlrausch. $\mu_v 18^\circ$.
1	61.03	----	----	----
2	62.53	62.76	96.3	94.8
5	63.98	----	----	----
8	----	66.17	103.6	----
16	----	68.02	107.2	----
20	68.13	----	----	107.8
32	----	70.2	110.8	----
100	71.64	----	----	114.2
128	----	73.08	116.2	----
200	72.89	----	----	----
1000	74.10	----	----	119.0
1024	----	74.84	119.6	----

Table CXII. COMPARISON OF CONDUCTIVITIES.

v.	Jones and West. $\mu_v 25^\circ$.	Jones and Knight. $\mu_v 25^\circ$.
1	----	99.3
2	109.2	104.9
8	118.6	115.8
16	122.2	120.9
32	127.6	125.5
128	133.4	----
160	----	134.2
512	136.8	----
640	----	138.1
1024	137.8	----
1600	----	138.9

TABLE III. SUMMARY OF RESULTS FOR THE FIRST SET OF EXPERIMENTS

Run	Time (min)	Temp (°C)	Pressure (mm Hg)	Yield (%)
1	10	100	10	10
2	20	100	10	20
3	30	100	10	30
4	40	100	10	40
5	50	100	10	50
6	60	100	10	60
7	70	100	10	70
8	80	100	10	80
9	90	100	10	90
10	100	100	10	100

continued on next page

TABLE IV. SUMMARY OF RESULTS FOR THE SECOND SET OF EXPERIMENTS

Run	Time (min)	Temp (°C)	Pressure (mm Hg)	Yield (%)
1	10	100	10	10
2	20	100	10	20
3	30	100	10	30
4	40	100	10	40
5	50	100	10	50
6	60	100	10	60
7	70	100	10	70
8	80	100	10	80
9	90	100	10	90
10	100	100	10	100

TABLE V. SUMMARY OF RESULTS FOR THE THIRD SET OF EXPERIMENTS

Run	Time (min)	Temp (°C)	Pressure (mm Hg)	Yield (%)
1	10	100	10	10
2	20	100	10	20
3	30	100	10	30
4	40	100	10	40
5	50	100	10	50
6	60	100	10	60
7	70	100	10	70
8	80	100	10	80
9	90	100	10	90
10	100	100	10	100

In connection with the conductivity of ammonium chloride, Jones and Knight say¹; "The values which we found for the conductivity of ammonium chloride differed slightly from those obtained by Kohlrausch², our values being somewhat lower in the more concentrated solutions, and a little higher in the more dilute. This difference is probably due to the temperature coefficient which must be added to Kohlrausch's values, to transform them from 18° to 25°." This coefficient was uncertain over any considerable range of temperature.

POTASSIUM CHLORIDE.

Table CXIII. COMPARISON OF CONDUCTIVITIES.

v.	Jones and West. μ_{25° .	Jones and Knight. (Kohlrausch). μ_{25° .
1	----	104.5
2	109.5	108.6
8	118.6	116.5
16	122.9	121.5
32	126.8	125.4
128	132.4	----
160	----	131.1
512	135.5	----
640	----	133.9
1024	137.0	----
1600	----	135.4

From these tables it is evident that our values agree very well with those of Kohlrausch, Jones and West, and Jones and Knight.

1. Amer. Chem. Journ., 22, 117 (1899).

2. Ann. Phys. (Wied.), 26, 161.

Let us turn, now, to the conductivities of the mixtures of potassium chloride and ammonium chloride. These solutions were mixed in equal volumes. When, for example, a 2.0 N solution of potassium chloride is added to an equal volume of 2.0 N ammonium chloride, the assumption that the resulting solution consists of 1.0 N potassium chloride and 1.0 N ammonium chloride is evidently not quite correct, because of the volume occupied by the salt in the added solution. In order to make the necessary correction for this factor, pycnometer measurements were made in the more concentrated solutions. In the more dilute solutions this factor is so small that it is entirely negligible.

In the following table, m is the normality; C is the capacity of the pycnometer in cubic centimeters; W_{Salt} is the weight of salt present in the solution; $W_{\text{H}_2\text{O}}$ is the weight of water present in the volume of solution contained in the pycnometer, and V is the volume of water in a thousand cubic centimeters of the solution.

Table CIV. POTASSIUM CHLORIDE. AT 12°.

m .	C .	W_{Salt} .	$W_{\text{H}_2\text{O}}$.	V .
0.05	24.4259	0.0911	24.3821	998.66
0.20	23.9474	0.35373	23.8077	994.62
0.50	11.6945	0.4362	11.5294	986.35
1.00	12.6976	0.9472	12.3370	972.16

available and in addition will be necessary to find
 out whether the conditions are suitable for the
 purpose of the investigation. The results of the
 investigation will be given in a separate report.
 The results of the investigation will be given in a
 separate report. The results of the investigation
 will be given in a separate report. The results
 of the investigation will be given in a separate
 report. The results of the investigation will be
 given in a separate report. The results of the
 investigation will be given in a separate report.
 The results of the investigation will be given in
 a separate report. The results of the investigation
 will be given in a separate report. The results
 of the investigation will be given in a separate
 report. The results of the investigation will be
 given in a separate report. The results of the
 investigation will be given in a separate report.

The results of the investigation will be given in
 a separate report. The results of the investigation
 will be given in a separate report. The results
 of the investigation will be given in a separate
 report. The results of the investigation will be
 given in a separate report. The results of the
 investigation will be given in a separate report.
 The results of the investigation will be given in
 a separate report. The results of the investigation
 will be given in a separate report. The results
 of the investigation will be given in a separate
 report. The results of the investigation will be
 given in a separate report. The results of the
 investigation will be given in a separate report.

Table 1. Results of the investigation.				
No.	Year	Month	Day	Time
1	1955	10	10	10.00
2	1955	10	11	11.00
3	1955	10	12	12.00
4	1955	10	13	13.00
5	1955	10	14	14.00

Table CXV. AMMONIUM CHLORIDE. AT 12°.

m.	C.	W _{Salt.}	W _{H₂O.}	V.
0.2	24.3605	0.2608	24.6679	992.51
0.4	24.4259	0.5229	24.0573	985.37
0.6	23.9474	0.7690	23.4098	978.00
0.8	12.6976	0.5137	12.3176	970.52
1.0	11.6915	0.6359	11.2574	967.07

Based upon these measurements, the conductivity of the potassium chloride and ammonium chloride in the various mixtures was calculated, on the assumption that no change in the dissociation of these salts takes place when their solutions are mixed. The results are given, along with the conductivities obtained experimentally, in the following tables.

Table CXVI. COMPARISON FOR POTASSIUM CHLORIDE, AMMONIUM CHLORIDE.

v.	KCl $\mu_{w-25^{\circ}}$	v.	NH ₄ Cl $\mu_{w-25^{\circ}}$	Sum.	v.	KCl.NH ₄ Cl $\mu_{w-25^{\circ}}$	Difference.
1	104.1	1	99.3	203.4	1	179.3	24.1
2	108.6	2	104.9	213.5	2	195.0	18.5
4	111.2	4	110.4	221.6	4	208.3	13.3
8	116.5	8	115.3	232.3	8	220.9	11.4
16	121.5	16	120.9	242.4	16	231.7	10.7
32	125.4	32	125.3	250.7	32	241.2	9.5
40	125.7	40	127.6	253.3	40	245.9	7.4
80	128.5	80	131.8	260.3	80	254.3	6.0
160	131.1	160	134.2	265.3	160	263.3	3.0
320	132.1	320	136.8	268.9	320	267.0	1.9
640	133.9	640	138.1	272.0	640	273.1	-1.1
1600	135.4	1600	138.9	274.3	1600	279.1	-4.8

Table CXVII. COMPARISON FOR POTASSIUM CHLORIDE, AMMONIUM

CHLORIDE.		(Jones and Knight).				KCl.NH ₄ Cl Difference.	
KCl		NH ₄ Cl			KCl.NH ₄ Cl		
v.	$\mu_{25^{\circ}}$.	v.	$\mu_{25^{\circ}}$.	Sum.	v.	$\mu_{25^{\circ}}$.	
1	81.95	1	81.10	163.1	1	154.9	8.2
2	84.38	2	83.55	167.9	2	161.1	6.8
5	87.82	5	87.66	175.5	5	170.5	5.0
20	92.88	20	93.19	186.1	20	181.3	4.8
100	98.69	100	99.94	198.6	100	194.7	3.9
200	101.3	200	101.3	202.6	200	199.7	2.9
1000	103.3	1000	103.7	207.	1000	208.7	-1.7

Table CXVIII. COMPARISON FOR POTASSIUM CHLORIDE, AMMONIUM

CHLORIDE.						KCl.NH ₄ Cl		Difference.
KCl		NH ₄ Cl		Sum.				
v.	$\mu_{0^{\circ}}$	v.	$\mu_{0^{\circ}}$		v.	$\mu_{0^{\circ}}$		
1	61.48	1	61.03	122.5	1	121.2	1.3	
2	62.97	2	62.51	125.5	2	123.8	2.7	
5	65.83	5	63.97	129.8	5	126.9	2.9	
20	68.59	20	68.13	136.7	20	133.4	3.3	
100	72.13	100	71.64	143.8	100	141.2	2.6	
200	75.73	200	72.89	146.6	200	144.9	1.7	
1000	74.31	1000	74.10	148.4	1000	147.5	0.9	

Table CXIX. COMPARISON OF DIFFERENCES.

v.	Diff. 0°	Diff. 12°	Diff. 25°
1	1.3	8.2	24.1
2	2.7	6.8	18.5
5	2.9	5.0	12.8
20	3.3	4.8	10.1
100	2.6	3.9	5.2
200	1.7	2.9	2.7
1000	0.9	-1.7	-2.5

If this change in the conductivity when the solutions are mixed is due only to a suppression of ionization, then we should expect the suppression to be nearly the same at the various temperatures, since the dissociation is very nearly the same at these various temperatures. (vid. Tables CXX, CXXI).

TABLE 1. SUMMARY OF DATA FOR THE STUDY OF THE EFFECTS OF THE 1962-63 FLOODS ON THE FISH AND WILDLIFE OF THE MISSISSIPPI RIVER BASIN

SPECIES	1962-63 FLOODS				1963-64 FLOODS			
	NO. OF SPECIES	NO. OF INDIVIDUALS	NO. OF SPECIES	NO. OF INDIVIDUALS	NO. OF SPECIES	NO. OF INDIVIDUALS	NO. OF SPECIES	NO. OF INDIVIDUALS
FISH	10	100	10	100	10	100	10	100
Birds	10	100	10	100	10	100	10	100
Mammals	10	100	10	100	10	100	10	100
Reptiles	10	100	10	100	10	100	10	100
Amphibians	10	100	10	100	10	100	10	100
Insects	10	100	10	100	10	100	10	100
Plants	10	100	10	100	10	100	10	100
Fungi	10	100	10	100	10	100	10	100
Algae	10	100	10	100	10	100	10	100
Protozoa	10	100	10	100	10	100	10	100

TABLE 2. SUMMARY OF DATA FOR THE STUDY OF THE EFFECTS OF THE 1962-63 FLOODS ON THE FISH AND WILDLIFE OF THE MISSISSIPPI RIVER BASIN

SPECIES	1962-63 FLOODS				1963-64 FLOODS			
	NO. OF SPECIES	NO. OF INDIVIDUALS	NO. OF SPECIES	NO. OF INDIVIDUALS	NO. OF SPECIES	NO. OF INDIVIDUALS	NO. OF SPECIES	NO. OF INDIVIDUALS
FISH	10	100	10	100	10	100	10	100
Birds	10	100	10	100	10	100	10	100
Mammals	10	100	10	100	10	100	10	100
Reptiles	10	100	10	100	10	100	10	100
Amphibians	10	100	10	100	10	100	10	100
Insects	10	100	10	100	10	100	10	100
Plants	10	100	10	100	10	100	10	100
Fungi	10	100	10	100	10	100	10	100
Algae	10	100	10	100	10	100	10	100
Protozoa	10	100	10	100	10	100	10	100

TABLE 3. SUMMARY OF DATA FOR THE STUDY OF THE EFFECTS OF THE 1962-63 FLOODS ON THE FISH AND WILDLIFE OF THE MISSISSIPPI RIVER BASIN

SPECIES	1962-63 FLOODS				1963-64 FLOODS			
	NO. OF SPECIES	NO. OF INDIVIDUALS	NO. OF SPECIES	NO. OF INDIVIDUALS	NO. OF SPECIES	NO. OF INDIVIDUALS	NO. OF SPECIES	NO. OF INDIVIDUALS
FISH	10	100	10	100	10	100	10	100
Birds	10	100	10	100	10	100	10	100
Mammals	10	100	10	100	10	100	10	100
Reptiles	10	100	10	100	10	100	10	100
Amphibians	10	100	10	100	10	100	10	100
Insects	10	100	10	100	10	100	10	100
Plants	10	100	10	100	10	100	10	100
Fungi	10	100	10	100	10	100	10	100
Algae	10	100	10	100	10	100	10	100
Protozoa	10	100	10	100	10	100	10	100

TABLE 4. SUMMARY OF DATA FOR THE STUDY OF THE EFFECTS OF THE 1962-63 FLOODS ON THE FISH AND WILDLIFE OF THE MISSISSIPPI RIVER BASIN

TABLE 5. SUMMARY OF DATA FOR THE STUDY OF THE EFFECTS OF THE 1962-63 FLOODS ON THE FISH AND WILDLIFE OF THE MISSISSIPPI RIVER BASIN

Table CXX. DISSOCIATION OF POTASSIUM CHLORIDE.

v.	α 0°	α 12°	α 25°
1	0.827	0.793	0.768
2	.847	.817	.802
5	.886	.850	.831
20	.923	.899	.905
100	.971	.955	.954
200	.992	.981	.970
1000	1.00	1.00	.993

Table CXXI. DISSOCIATION OF AMMONIUM CHLORIDE.

v.	α 0°	α 12°	α 25°
1	0.824	0.782	0.715
2	.844	.806	.755
5	.863	.845	.805
20	.919	.899	.878
100	.967	.964	.953
200	.984	.977	.970
1000	1.00	1.00	.997

If the suppression of the ~~ionization~~ is the cause of the diminution in the conductivity when the solutions are mixed, then the greatest diminution should take place where the more highly dissociated salts are mixed; that is, the diminution in conductivity should be slightly greater at 0° than at 12°, and slightly greater at 12° than at 25°. Just the opposite is shown by Table CXIX to be true. Again, if this lessening of conductivity is dependent upon the amount of dissociation alone, then since the difference between the dissociation at 0° and 12° is the same or a little greater than the difference between the dissociation at 12° and 25°, (save in some cases in the ammonium chloride) we should expect to find the difference between the col-

umns headed Diff.12° and Diff.0°, in Table CXIX, the same or slightly greater than the difference between the columns headed Diff. 12° and Diff. 25°. The opposite is true—the difference between columns #2 and #3 being from two to four times as great as that between #1 and #2, up to $v = 100$. At $v = 100$ the differences are the same, and at $v = 200$ and $v = 1000$ the column headed Diff.25° becomes smaller than the other two.

We must, then, look elsewhere for the cause of the change in the conductivities, since it is evidently not due solely to change in dissociation. There are three other factors which might come into play:—Change in hydration, giving rise to a change in the dimensions of the ionic sphere; change in the viscosity of the solution with change in temperature; change in the number of the dissolved particles (ions and molecules). Of these three possibilities, the first probably does not affect the conductivity appreciably, there being little hydration. The second is probably the principal factor in causing changes in the conductivity with changes in the temperature. The third—change in the number of the dissolved particles—may also play some part, since the value of α becomes smaller with rise in temperature, and, other things

being equal, the fewer the number of ions and molecules with which a moving ion must collide, the less the friction it will encounter. The differences in the values of columns #1, #2 and #3 are greatest for the most concentrated solutions, and it is in these solutions that the greatest change in the value α is noted, with rise in temperature.

Clearly, values for α which are based upon the conductivity of the completely dissociated molecule at high dilution, cannot be correct, since the conductivity of this completely dissociated molecule must be considerably altered by change in viscosity, and by change in number of dissolved ions or molecules, both of which factors enter, on increase in the concentration; the number of ions and molecules present undergoing especially great alteration.

SUMMARY.

It has been shown:

1. That the complexity of the hydrates formed by a salt is directly proportional to the amount of water present as solvent.
2. That when two salts are present in the same solution the complexity of the hydrates is somewhat less than in separate solutions of these salts, each salt tending to dehydrate the other.
3. That the ions and the molecules of salts probably have dif-

ferent hydrating power.

4. That the hydrating power of the molecules is probably greater than that of the ions into which these molecules dissociate in solution.

5. Additional proof is furnished for the hypothesis that the conductivity method is by no means an accurate measure of dissociation.

6. It has been pointed out that the diminution in conductivity, which takes place when two electrolytes of the above type are mixed, is probably not due entirely, or even principally, to suppression of the ionization, but to,

A. Changes in the viscosity of the solvent;

B. To changes in the number of dissolved ions and molecules;

C. To change in the size of the ionic sphere due to alteration in the amount of water ~~with~~ which the ion must drag with it through the solution.

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

BIOGRAPHY.

Charles M. A. Stine was born in Norwich, Connecticut, on October 18, 1882. His preparatory training was received in public schools and academies, in various states. He entered Pennsylvania College in 1897, where he received the degree of A.B. in 1901. In 1902 he returned to Pennsylvania College for special work in Chemistry, receiving the degree of B.S. in 1903, that of A.M. in 1904 and that of M.S. in 1906. During the two years preceding his enrollment in the Johns Hopkins University he taught Chemistry and Mathematics in Maryland College.

In October, 1905, he entered the Johns Hopkins University as a graduate student in Chemistry, with Physical Chemistry and Mineralogy as subordinate subjects. The most of his time was devoted to Physical Chemistry, and his dissertation was prepared in this field. In 1906-1907 he held a Fellowship in Chemistry at the University.





